

# Datasheet: AHP3057 BATCH NUMBER 160823

| Description:  | RABBIT ANTI TRI-METHYL-HISTONE H3 (Lys36) |  |  |
|---------------|---|--|--|
| Specificity:  | TRI-METHYL-HISTONE H3 (Lys36)             |  |  |
| Format:       | Purified                                  |  |  |
| Product Type: | Polyclonal Antibody                       |  |  |
| Isotype:      | Polyclonal IgG                            |  |  |
| Quantity:     | 0.1 mg                                    |  |  |
|               |   |  |  |

## **Product Details**

### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="https://www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

|                     | Yes | No | Not Determined | Suggested Dilution  |
|---------------------|-----|----|----------------|---------------------|
| Western Blotting    | •   |    |                | 0.5 μg/ml - 2 μg/ml |
| Chromatin           | _   |    |                | 10 µg per ChIP      |
| Immunoprecipitation | _   |    |                | το μg per Chir      |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

| Target Species              | Human   |
|-----------------------------|---|
| Species Cross<br>Reactivity | Reacts with: Mouse Based on sequence similarity, is expected to react with:Broad <b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. |
| Product Form                | Purified IgG - liquid   |
| Antiserum Preparatio        | n Antiserum to tri-methyl-histone H3 (Lys36) was raised by repeated immunization of rabbits with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography on Protein A.   |
| Buffer Solution             | Phosphate buffered saline   |

**Preservative** 0.035% Sodium Azide (NaN<sub>3</sub>) **Stabilisers** 30% Glycerol **Carrier Free** Yes **Immunogen** A peptide containing tri-methylated lysine 36 of human histone H3 **External Database UniProt:** Links P68431 Related reagents **Entrez Gene:** 8968 HIST1H3F Related reagents **Synonyms** H3FA, H3FB, H3FC, H3FD, H3FF, H3FH, H3FI, H3FJ, H3FK, H3FL **Specificity** Rabbit anti Human tri-methyl-histone H3 (Lys36) antibody recognizes histone H3 when tri-methylated at lysine 36. Histone H3 is one of the four core histones that make up the nucleosome core particle. Nucleosomes are the smallest subunit of chromatin and are made up of 146 bp of DNA wrapped around an octamer comprised of pairs of the four core histones (H2A, H2B, H3, and H4) (Smith, 1991). Histones can be mono-, di- or tri-methylated by histone methyltransferases. Depending on which amino acid residues are methylated, methylation of histones may increase or decrease the transcription of genes. Methylation events that weaken the binding between histone tails and DNA lead to increased transcription because they make the DNA more accessible to transcription factor proteins and RNA polymerase. Methylation of histones is therefore crucial for the regulation of gene expression (Gupta et al. 2010). Tri-methylation of histone H3 at lysine 36 has been shown to lead to transcriptional activation (Xu et al. 2008). Wide species cross-reactivity is expected from Rabbit anti Human tri-methyl-histone H3 (Lys36) antibody based on sequence. Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch **Health And Safety** Material Safety Datasheet documentation #10049 available at: Information https://www.bio-rad-antibodies.com/SDS/AHP3057

10049

# **Related Products**

## **Recommended Secondary Antibodies**

Sheep Anti Rabbit IgG (STAR34...) FITC

Sheep Anti Rabbit IgG (STAR35...) RPE

Goat Anti Rabbit IgG (H/L) (STAR124...) HRP

Goat Anti Rabbit IgG (Fc) (STAR121...) Biotin, FITC, HRP

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M394884:220218'

#### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint