

Datasheet: AHP3076

| | |
|----------------------|-------------------------|
| Description: | RABBIT ANTI HUMAN HMGN2 |
| Specificity: | HMGN2 |
| Format: | Purified |
| Product Type: | Polyclonal Antibody |
| Isotype: | Polyclonal IgG |
| Quantity: | 0.1 ml |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|------------------|-----|----|----------------|--------------------|
| Western Blotting | ▪ | | | 1/500 - 1/2000 |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

| | |
|------------------------------|--|
| Target Species | Human |
| Product Form | Purified IgG - liquid |
| Antiserum Preparation | Antiserum to human HMGN2 was raised by repeated immunization of rabbits with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography. |
| Buffer Solution | Phosphate buffered saline |
| Preservative | 0.035% Sodium Azide (NaN ₃) |
| Stabilisers | 30% Glycerol |
| Carrier Free | Yes |
| Immunogen | A peptide within the N-terminal region of human HMGN2 |

External Database Links

UniProt:
[P05204](https://www.uniprot.org/entry/P05204) [Related reagents](#)

Entrez Gene:[3151](#) HMGN2 [Related reagents](#)**Synonyms**

HMG17

Specificity

Rabbit anti Human HMGN2 antibody recognizes high mobility group nucleosome-binding domain-containing protein 2, also known as non-histone chromosomal protein HMG-17.

HMGN2 is a 9 kDa member of the HMG class of chromosomal proteins, the most abundant non-histones present in the nuclei of eukaryotic cells ([Hock et al. 2007](#)). Chromatin regions with high levels of HMGN2 are enriched in transcriptionally active genes and in acetylated histones. HMGN2 associates with actively transcribed genes. HMGN2 also binds to the inner side of the nucleosomal DNA, thereby regulating the interaction between the DNA and the nucleosome. HMGN2 may also play a role in maintaining transcribable genes in a unique chromatin conformation ([Garza-Manero et al. 2019](#)).

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10049 available at: <https://www.bio-rad-antibodies.com/SDS/AHP3076>
10049

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...) [FITC](#)Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)Sheep Anti Rabbit IgG (STAR35...) [RPE](#)Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)**North & South** Tel: +1 800 265 7376**America** Fax: +1 919 878 3751Email: antibody_sales_us@bio-rad.com**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M394899:220218'

