



Datasheet: AHP318

BATCH NUMBER 161516

Description:	RABBIT ANTI DNA PKcs
Specificity:	DNA PKcs
Format:	Serum
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin (1)	▪			1/100
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting	▪			1/2000
Gel Super Shift Assays	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)This product requires heat mediated treatment of paraffin sections. Bio-Rad recommend the use of antigen unmasking fluid for this purpose. A two hour incubation with AHP318 is recommended.

Target Species

Human

Species Cross Reactivity

Reacts with: Hamster, Mouse, Rat

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Serum - liquid

Antiserum Preparation	Antisera to DNA-PKcs were raised by repeated immunisation of rabbit with highly purified antigen. Whole serum is supplied.
Preservative Stabilisers	<0.1% Sodium Azide (NaN ₃)
Immunogen	Human DNA-PKcs fragment.
External Database Links	<p>UniProt: P78527 Related reagents</p> <p>Entrez Gene: 5591 PRKDC Related reagents</p>
Synonyms	HYRC, HYRC1
RRID	AB_322027
Specificity	Rabbit anti DNA PKcs antibody recognizes the human DNA-dependent protein kinase catalytic subunit, also known as DNA-PKcs, DNPk1 or p460. DNA PKcs is a 4128 amino acid ~460 kDa nuclear protein encoded by the PRKDC gene, acting as a molecular sensor for DNA damage. Mutation of the PRKDC gene can result in Immunodeficiency 26 with or without neurologic abnormalities (IMD26) (van der Burg et al. 2009).
References	<ol style="list-style-type: none"> 1. Finnie, N.J. <i>et al.</i> (1995) DNA-dependent protein kinase activity is absent in xrs-6 cells: implications for site-specific recombination and DNA double-strand break repair. Proc Natl Acad Sci U S A. 92 (1): 320-4. 2. Jayaram, S. <i>et al.</i> (2008) E1B 55k-independent dissociation of the DNA ligase IV/XRCC4 complex by E4 34k during adenovirus infection. Virology.382: 163-70. 3. Hoek, M. <i>et al.</i> (2011) An analysis of CAF-1-interacting proteins reveals dynamic and direct interactions with the KU complex and 14-3-3 proteins. J Biol Chem. 286 (12): 10876-87. 4. Frasca, D. <i>et al.</i> (2001) Role of DNA-dependent protein kinase in recognition of radiation-induced DNA damage in human peripheral blood mononuclear cells. Int Immunol. 13: 791-7. 5. Jayaram, S. <i>et al.</i> (2008) Loss of DNA ligase IV prevents recognition of DNA by double-strand break repair proteins XRCC4 and XLF. Nucleic Acids Res. 36: 5773-86. 6. Koike, M. <i>et al.</i> (1999) Differential subcellular localization of DNA-dependent protein kinase components Ku and DNA-PKcs during mitosis. J Cell Sci. 112: 4031-9. 7. Yee, M.C. <i>et al.</i> (2005) A cell-permeable, activity-based probe for protein and lipid kinases. J Biol Chem. 280: 29053-9. 8. Zhao, H.J. <i>et al.</i> (2000) DNA-dependent protein kinase activity correlates with Ku70 expression and radiation sensitivity in esophageal cancer cell lines. Clin Cancer Res. 6: 1073-8. 9. Sirzen, F. <i>et al.</i> (1999) DNA-dependent protein kinase content and activity in lung carcinoma cell lines: correlation with intrinsic radiosensitivity. Eur J Cancer. 35: 111-6. 10. Frasca, D. <i>et al.</i> (2002) The DNA repair protein ku is involved in gp130-mediated

signal transduction events in PBMC from young but not from elderly subjects. [Exp Gerontol. 37: 321-8.](#)

11. Scarpaci, S. *et al.* (2003) DNA damage recognition and repair capacities in human naïve and memory T cells from peripheral blood of young and elderly subjects. [Mech Ageing Dev. 124: 517-24.](#)

12. Kamachi, M. *et al.* (2002) Human autoimmune sera as molecular probes for the identification of an autoantigen kinase signaling pathway. [J Exp Med. 196: 1213-25.](#)

13. Gaymes, T.J. *et al.* (2006) Histone deacetylase inhibitors (HDI) cause DNA damage in leukemia cells: a mechanism for leukemia-specific HDI-dependent apoptosis? [Mol Cancer Res. 4 \(8\): 563-73.](#)

14. Koike, M. *et al.* (1999) Differential subcellular localization of DNA-dependent protein kinase components Ku and DNA-PKcs during mitosis. [J Cell Sci. 112 \(Pt 22\): 4031-9.](#)

15. Nilsson, A. *et al.* (1999) Cell cycle-dependent regulation of the DNA-dependent protein kinase. [Cell Prolif. 32 \(4\): 239-48.](#)

16. Yumoto Y *et al.* (1998) High mobility group proteins 1 and 2 can function as DNA-binding regulatory components for DNA-dependent protein kinase *in vitro*. [J Biochem. 124 \(3\): 519-27.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10081 available at: <https://www.bio-rad-antibodies.com/SDS/AHP318>
10081

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)

Sheep Anti Rabbit IgG (STAR35...) [RPE](#)

Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)

Recommended Useful Reagents

[TidyBlot WESTERN BLOT DETECTION REAGENT:HRP \(STAR209P\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M389221:210806'

