

## Datasheet: AHP340B

**BATCH NUMBER 144550**

<b>Description:</b>	RABBIT ANTI HUMAN RANTES:Biotin
<b>Specificity:</b>	RANTES
<b>Other names:</b>	CCL5
<b>Format:</b>	Biotin
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	50 µg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			■	
Immunohistology - Frozen			■	
Immunohistology - Paraffin	■			
ELISA	■			0.25 - 1.0ug/ml
Immunoprecipitation			■	
Western Blotting	■			0.1 - 0.2ug/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Human
<b>Product Form</b>	Purified IgG conjugated to Biotin - lyophilized
<b>Reconstitution</b>	<p>Reconstitute with 0.5 ml sterile PBS containing 0.1% Bovine Serum Albumin. Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution.</p> <p>For long term storage the addition of 0.09% sodium azide is recommended.</p>

**Antiserum Preparation** Antisera to human RANTES were raised by repeated immunisations of rabbits with highly purified antigen. Purified IgG prepared by affinity chromatography.

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	None present
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml after reconstitution.
Immunogen	<a href="#">Recombinant Human RANTES</a>
External Database Links	<p><b>UniProt:</b></p> <p><a href="#">P13501</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b></p> <p><a href="#">6352</a>    CCL5    <a href="#">Related reagents</a></p>
Synonyms	D17S136E, SCYA5
RRID	AB_2244130
Specificity	<p><b>Rabbit anti Human RANTES antibody</b> recognizes human RANTES (regulated upon activation, normal T cell expressed and secreted), otherwise known as CCL5, a 68 amino acid CC- chemokine involved in T cell activation and cell recruitment, expressed by cytotoxic (CD8+) T cells (CTLs), macrophages, keratinocytes and endothelial cells.</p> <p>RANTES acts as a chemoattractant for monocytes, CD4+/CD45RO+ memory T cells, basophils and eosinophils, thorough binding to chemokine (C-C motif) receptor 1 (CCR1), CCR3 and CCR5. Both CCR3 and CCR5 are known co-receptors for the entry of HIV viruses into target cells and the binding of RANTES to CCR3 on CTLs has been shown to enhance HIV-1-specific cytotoxicity, whilst binding to CCR5 inhibits membrane fusion and hence HIV entry.</p>
ELISA	This biotinylated human RANTES antibody may be used in a direct ELISA or as the detection reagent in a sandwich ELISA with our <a href="#">purified human RANTES antibody</a> (AHP340G) as the capture reagent and <a href="#">recombinant human RANTES</a> (PHP062) as the standard.
Western Blotting	This antibody may be used in Western Blotting under either reducing or non-reducing conditions with <a href="#">recombinant human RANTES</a> (PHP062) as the positive control.
References	<ol style="list-style-type: none"> <li>Appay, V. <i>et al.</i> (2000) RANTES activates antigen-specific cytotoxic T lymphocytes in a mitogen-like manner through cell surface aggregation. <a href="#">Int Immunol. 12 (8): 1173-82.</a></li> <li>Dragic, T. <i>et al.</i> (1996) HIV-1 entry into CD4+ cells is mediated by the chemokine receptor CC-CKR-5. <a href="#">Nature. 381 (6584): 667-73.</a></li> <li>Cheng, H. <i>et al.</i> (2016) TWEAK/Fn14 activation induces keratinocyte proliferation under psoriatic inflammation. <a href="#">Exp Dermatol. 25 (1): 32-7.</a></li> </ol>
Storage	Prior to reconstitution store at +4°C.

After reconstitution store at -20°C.

Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

<b>Guarantee</b>	6 months from date of reconstitution
------------------	--------------------------------------

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10162 available at: <a href="https://www.bio-rad-antibodies.com/SDS/AHP340B">https://www.bio-rad-antibodies.com/SDS/AHP340B</a> 10162
--------------------------------------	--

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)  
'M344725:190118'

**Printed on 28 May 2025**

---

© 2025 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)