

Datasheet: AHP423 BATCH NUMBER 071212

Description:	RABBIT ANTI SHEEP INTERLEUKIN-1 BETA
Specificity:	IL-1 BETA
Format:	Serum
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				
Immunohistology - Frozen				
Immunohistology - Paraffin				
ELISA				1/500
Immunoprecipitation			•	
Western Blotting				

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Sheep
Species Cross Reactivity	Reacts with: Bovine
. Cood and a second	N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
Product Form	Serum - liquid
Antiserum Prepara	tion Antisera to ovine IL-1 beta were raised by repeated immunisation of rabbits with highly purified antigen.

Pre	ser	vati	ve
Sta	bili	sers	

0.1% Sodium Azide (NaN₃)

Immunogen

Recombinant IL-1 beta.

External Database Links

UniProt:

P21621 Related reagents

Entrez Gene:

443539 IL-1B Related reagents

Specificity

Rabbit anti Sheep interleukin 1β antibody recognizes ovine interleukin-1 beta (IL-1 beta). No cross-reactivity is seen with ovine IL-6, IL-8, MCP-1 or TNF alpha.

Rabbit anti Sheep interleukin 1β antibody neutralizes the bioactivity of ovine IL-1 beta. Removal of sodium azide is recommended prior to use in functional assays.

References

- 1. Rothel, J.S. *et al.* (1997) Analysis of ovine IL-1 beta production *in vivo* and *in vitro* by enzyme immunoassay and immunohistochemistry. <u>Vet. Immunol. Immunopathol. 57:</u> 267-278.
- 2. Bannerman, D.D. *et al.* (2004) *Escherichia coli* and *Staphylococcus aureus* elicit differential innate immune responses following intramammary infection. <u>Clin Diagn Lab Immunol.</u> 11: 463-72.
- 3. Dernfalk, J. *et al.* (2007) The xMAP technique can be used for detection of the inflammatory cytokines IL-1beta, IL-6 and TNF-alpha in bovine samples. <u>Vet Immunol Immunopathol. 118: 40-9.</u>
- 4. Moyes, K.M. *et al.* (2009) Identification of potential markers in blood for the development of subclinical and clinical mastitis in dairy cattle at parturition and during early lactation. J Dairy Sci. 92: 5419-28.
- 5. Allen, J.S. *et al.* (2009) Plasmacytoid dendritic cells are proportionally expanded at diagnosis of type 1 diabetes and enhance islet autoantigen presentation to T-cells through immune complex capture. <u>Diabetes</u>. 2009 Jan;58(1):138-45.
- 6. Vordermeier, M. *et al.* (2012) Cytokine responses of Holstein and Sahiwal zebu derived monocytes after mycobacterial infection. Trop Anim Health Prod. 44: 651-5.
- 7. Xu, A. *et al.* (2015) The Ovine Fetal and Placental Inflammatory Response to Umbilical Cord Occlusions With Worsening Acidosis. Reprod Sci. pii: 1933719115580994.
- 8. Bagu, E.T. *et al.* (2010) Post-natal changes in testicular concentrations of interleukin-1 alpha and beta and interleukin-6 during sexual maturation in bulls. <u>Reprod Domest Anim.</u> 45 (2): 336-41.
- 9. Atik, A. *et al.* (2012) Long-term pulmonary effects of intrauterine exposure to endotoxin following preterm birth in sheep. Reprod Sci. 19 (12): 1352-64.
- 10. Silvestre, F.T. *et al.* (2011) Effects of differential supplementation of fatty acids during the peripartum and breeding periods of Holstein cows: II. Neutrophil fatty acids and function, and acute phase proteins. <u>J Dairy Sci. 94 (5): 2285-301.</u>

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation available at: https://www.bio-rad-antibodies.com/SDS/AHP423 Material Safety Datasheet Documentation #10342 available at: https://www.bio-rad-antibodies.com/uploads/MSDS/10342.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (H/L) (STAR124...) HRP Sheep Anti Rabbit IgG (STAR35...)

Goat Anti Rabbit IgG (Fc) (STAR121...) Biotin, FITC, HRP

Recommended Useful Reagents

TidyBlot WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

Worldwide

North & South Tel: +1 800 265 7376 America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M279405:160115'

Printed on 20 Mar 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint