

Datasheet: AHP423 BATCH NUMBER 147881

Description:	RABBIT ANTI SHEEP INTERLEUKIN-1 BETA
Specificity:	IL-1 BETA
Format:	Serum
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			•	
Immunohistology - Frozen			•	
Immunohistology - Paraffin			•	
ELISA	•			1/500
Immunoprecipitation			•	
Western Blotting				

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Sheep
Species Cross Reactivity	Reacts with: Bovine Based on sequence similarity, is expected to react with:Goat N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
Product Form	Serum - liquid
Antiserum Prepara	tion Antisera to ovine IL-1 beta were raised by repeated immunisation of rabbits with highly purified antigen.

Immunogen Recombinant IL-1 beta.	
Preservative Stabilisers 0.09% Sodium Azide (NaN ₃)	

P21621 Related reagents

Entrez Gene:

443539 IL-1B Related reagents

RRID

AB 322127

Specificity

Rabbit anti Sheep interleukin 1 β antibody recognizes ovine interleukin-1 beta (IL-1 beta). No cross-reactivity is seen with ovine IL-6, IL-8, MCP-1 or TNF alpha.

Rabbit anti Sheep interleukin 1β antibody neutralizes the bioactivity of ovine IL-1 beta. Removal of sodium azide is recommended prior to use in functional assays.

References

- 1. Rothel, J.S. *et al.* (1997) Analysis of ovine IL-1 beta production *in vivo* and *in vitro* by enzyme immunoassay and immunohistochemistry. <u>Vet Immunol Immunopathol. 57 (3-4):</u> 267-78.
- 2. Bannerman, D.D. *et al.* (2004) *Escherichia coli* and *Staphylococcus aureus* elicit differential innate immune responses following intramammary infection. <u>Clin Diagn Lab Immunol.</u> 11: 463-72.
- 3. Dernfalk, J. *et al.* (2007) The xMAP technique can be used for detection of the inflammatory cytokines IL-1beta, IL-6 and TNF-alpha in bovine samples. <u>Vet Immunol Immunopathol</u>. 118: 40-9.
- 4. Moyes, K.M. *et al.* (2009) Identification of potential markers in blood for the development of subclinical and clinical mastitis in dairy cattle at parturition and during early lactation. J Dairy Sci. 92: 5419-28.
- 5. Silvestre, F.T. *et al.* (2011) Effects of differential supplementation of fatty acids during the peripartum and breeding periods of Holstein cows: II. Neutrophil fatty acids and function, and acute phase proteins. <u>J Dairy Sci. 94 (5): 2285-301.</u>
- 6. Vordermeier, M. *et al.* (2012) Cytokine responses of Holstein and Sahiwal zebu derived monocytes after mycobacterial infection. <u>Trop Anim Health Prod. 44: 651-5.</u>
- 7. Xu, A. *et al.* (2015) The Ovine Fetal and Placental Inflammatory Response to Umbilical Cord Occlusions With Worsening Acidosis. Reprod Sci. 22 (11): 1409-20.
- 8. Bagu, E.T. *et al.* (2010) Post-natal changes in testicular concentrations of interleukin-1 alpha and beta and interleukin-6 during sexual maturation in bulls. <u>Reprod Domest Anim.</u> 45 (2): 336-41.
- 9. Atik, A. *et al.* (2012) Long-term pulmonary effects of intrauterine exposure to endotoxin following preterm birth in sheep. Reprod Sci. 19 (12): 1352-64.
- 10. Cortes, M. *et al.* (2016) RNAseq profiling of primary microglia and astrocyte cultures in near-term ovine fetus: a glial *in vivo-in vitro* multi-hit paradigm in large mammalian brain Journal of Neuroscience Methods. Nov 14 [Epub ahead of print]
- 11. Doull, L. et al. (2015) Late production of CXCL8 in ruminant oro-nasal turbinate cells in

response to *Chlamydia* abortus infection. <u>Vet Immunol Immunopathol. 168 (1-2): 97-102.</u> 12. Cao, M. *et al.* (2019) α7 Nicotinic Acetylcholine Receptor Signaling Modulates Ovine Fetal Brain Astrocytes Transcriptome in Response to Endotoxin. Front Immunol. 10: 1063.

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (Fc) (STAR121...) Biotin, FITC, HRP

Sheep Anti Rabbit IgG (STAR35...) RPE

Goat Anti Rabbit IgG (H/L) (STAR124...) HRP

Email: antibody_sales_us@bio-rad.com

Recommended Useful Reagents

TidyBlot WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

North & South Tel: +1 800 265 7376

America

Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M364212:200529'

Printed on 15 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint