

# Datasheet: AHP455G

Description:	RABBIT ANTI HUMAN INTERLEUKIN-17
Specificity:	IL-17
Other names:	IL-17A
Format:	Purified
Product Type:	Polyclonal Antibody
lsotype:	Polyclonal IgG
Quantity:	0.1 mg

# **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further						
		information. For general protocol recommendations, please visit www.bio-					
	rad-antibodies.com/proto						
		Yes	No	Not Determined	Suggested Dilution		
	Flow Cytometry			•			
	Immunohistology - Frozen			•			
	Immunohistology - Paraffin (1)	•			0.25ug/ml		
	ELISA	-			0.5 - 2.0ug/ml		
	Immunoprecipitation			•			
	Western Blotting	-			0.1 - 0.2ug/ml		
	Immunofluorescence	-					
	Functional Assays (2)	-			0.93 - 1.3ug/ml		
	<ul> <li>Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.</li> <li>(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.</li> <li>(2) To yield one-half maximal inhibition [ND50] of the biological activity of hIL-17A (50.0 ng/ml).</li> </ul>						
Target Species	Human						
Product Form	Purified IgG - lyophilized						
Reconstitution	Reconstitute with 0.1ml c Care should be taken du bottom of the vial. Bio-Ra	ring reco	nstitution				

For long term storage the addition of 0.09% sodium azide is recommended. N.B. For functional studies do not add sodium azide

Antiserum Preparation Antisera to human IL-17 were raised by repeated immunisation of rabbits with highly purified antigen. Purified IgG prepared by antigen affinity chromatography

		0 1 9		
Buffer Solution	Phosphate buffered saline			
Preservative Stabilisers	None present			
Carrier Free	Yes			
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml after reconstitution			
Immunogen	Recombinant human IL-17			
External Database Links	UniProt: <u>Q16552</u> <u>Related reagents</u> Entrez Gene: <u>3605</u> IL17A <u>Related reagents</u>			
Synonyms	CTLA8, IL17			
RRID	AB_2124883			
Specificity	<b>Rabbit anti Human Interleukin-17 antibody</b> recognizes human IL-17 (Interleukin- 17/17A), also known as Cytotoxic T-lymphocyte-associated antigen 8 or CTLA-8. Interleukin-17 is a homodimer of two 132 amino acid chains resulting oin a ~32 kDa disulfide-linked cytokine belonging to the IL-17 family (classified as IL-17A to IL-17F) which signals through the IL-17 receptor (IL-17R/CDw217). IL-17 is produced primarily by activated CD4+ T cells and induces a variety of target cells to secrete inflammatory cytokines, including IL-1beta, IL-6, TNF- $\alpha$ , IFN-γ, and granulocyte colony-stimulating factor, as well as synergizing with other inflammatory cytokines and agonists. IL-17 has roles in bone metabolism and rheumatoid arthritis, in which levels of IL-17 are significantly increased and may contribute to bone destruction (Ziolkowska <i>et al.</i> 2000).			
ELISA	This purified human IL-17 antibody may be used in an ELISA <u>IL-17</u> (PHP098) as the standard.	with <u>recombinant human</u>		
Western Blotting	This antibody may be used in Western Blotting under either re conditions with <u>recombinant human IL-17</u> (PHP098) as the pos			
References	1. Parsonage, G. <i>et al.</i> (2008) Prolonged, granulocyte-macrop	hage colony-stimulating		

	factor-dependent, neutrophil survival following rheumatoid synd IL-17 and TNFalpha. <u>Arthritis Res Ther. 10: R47.</u>	ovial fibroblast activation by
Further Reading	1. Gaffen, S.L. (2004) Biology of recently discovered cytokines inflammatory cytokine with roles in bone biology and arthritis. <u>A</u> 240-7.	
Storage	This product is shipped at ambient temperature.	
	Prior to reconstitution store at -20°C.	
	After reconstitution store at -20°C.	
	This product should be stored undiluted. Storage in frost-free f recommended. Avoid repeated freezing and thawing as this ma Should this product contain a precipitate we recommend micro	ay denature the antibody.
Guarantee	12 months from date of despatch	
Health And Safety Information	Material Safety Datasheet documentation #10294 available at: https://www.bio-rad-antibodies.com/SDS/AHP455G 10294	
Regulatory	For research purposes only	

## **Related Products**

### **Recommended Secondary Antibodies**

Goat Anti Rabbit IgG (H/L) (STAR124...) <u>HRP</u> Sheep Anti Rabbit IgG (STAR35...) <u>RPE</u> Goat Anti Rabbit IgG (Fc) (STAR121...) <u>Biotin</u>, <u>FITC</u>, <u>HRP</u> **Recommended Positive Controls** 

RECOMBINANT HUMAN INTERLEUKIN-17 (PHP098)

### **Recommended Useful Reagents**

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A) TidyBlot WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel:

 America
 Fax: +1 919 878 3751
 Fax

 Email: antibody\_sales\_us@bio-rad.com
 Email

Tel: +44 (0)1865 852 700 **Europe** Fax: +44 (0)1865 852 739 Email: antibody\_sales\_uk@bio-rad.com Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody\_sales\_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M439220:250523'

#### Printed on 23 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint