Datasheet: AHP500

BATCH NUMBER 156373

Description:	SHEEP ANTI HUMAN TGN46
Specificity:	TGN46
Other names:	TGOLN2
Format:	Serum
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen	▪			1/50 - 1/100
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting	▪			1/500 - 1/000
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species

Human

Species Cross Reactivity

Reacts with: Primate

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Serum, diluted - liquid

Antiserum Preparation

Antisera to human TGN46 were raised by repeated immunisation of sheep with highly

	purified antigen.
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN_3) 1% Bovine Serum Albumin 25% Glycerol
Immunogen	Recombinant human TGN46.
External Database Links	**UniProt:** O43493 Related reagents **Entrez Gene:** 10618 TGOLN2 Related reagents
Synonyms	TGN46, TGN51
RRID	AB_324049
Specificity	**Sheep anti Human TGN46 antibody** recognizes Trans-Golgi network integral membrane protein 2 (TGOLN2), also known as TGN38 homolog, TGN46, TGN48 or Trans-Golgi network protein TGN51. TGN46 is a 437 amino acid glycoprotein localized to the Trans-Golgi network. TGN46 has been reported as being the best available marker for human trans-Golgi network. TGN46 is a heavily glycosylated protein of around 110-120 kDa. Multiple isoforms of TGN46 are generated by alternative splicing difffering in sequence at the C-terminal protion. Sheep anti Human TGN46 antibody is expeced to recognize all identified isoforms.
Immunohistology	Fixation with 3% paraformaldehyde or methanol/acetone is recommended.
References	1. Prescott AR *et al.* (1997) Distinct compartmentalization of TGN46 and beta 1,4-galactosyltransferase in HeLa cells. Eur J Cell Biol. 72 (3): 238-46. 2. Vuillier, F. *et al.* (2005) Lower levels of surface B-cell-receptor expression in chronic lymphocytic leukemia are associated with glycosylation and folding defects of the mu and CD79a chains. Blood. 105 (7): 2933-40. 3. Edwards, T.L. *et al.* (2009) Endogenous spartin (SPG20) is recruited to endosomes and lipid droplets and interacts with the ubiquitin E3 ligases AIP4 and AIP5. Biochem J. 423 (1): 31-9. 4. Roberts, R.C. *et al.* (2010) Mistargeting of SH3TC2 away from the recycling endosome causes Charcot-Marie-Tooth disease type 4C. Hum Mol Genet.19: 1009-18. 5. Hauser, H. *et al.* (2010) HIV-1 Vpu and HIV-2 Env counteract BST-2/tetherin by sequestration in a perinuclear compartment. Retrovirology. 7: 51. 6. Oliver, S.L. *et al.* (2011) Mutagenesis of varicella-zoster virus glycoprotein I (gI) identifies a cysteine residue critical for gE/gI heterodimer formation, gI structure, and

virulence in skin cells. J Virol. 85 (9): 4095-110.
7. Berarducci, B. *et al.* (2006) Essential functions of the unique N-terminal region of the varicella-zoster virus glycoprotein E ectodomain in viral replication and in the pathogenesis of skin infection. J Virol. 80: 9481-96.
8. Sadaoka, T. *et al.* (2010) Characterization of the varicella-zoster virus ORF50 gene, which encodes glycoprotein M. J Virol. 84: 3488-502.
9. Vleck, S.E. *et al.* (2010) Anti-glycoprotein H antibody impairs the pathogenicity of varicella-zoster virus in skin xenografts in the SCID mouse model. J Virol. 84: 141-52.
10. Esk, C. *et al.* (2010) The clathrin heavy chain isoform CHC22 functions in a novel endosomal sorting step. J Cell Biol. 188: 131-44.
11. Cheng, S.B. *et al.* (2011) Down-modulation of the G-protein-coupled Estrogen Receptor, GPER, from the Cell Surface Occurs via a trans-Golgi-Proteasome Pathway. J Biol Chem. 286: 22441-55.
12. Fairn, G.D. *et al.* (2011) High-resolution mapping reveals topologically distinct cellular pools of phosphatidylserine. J Cell Biol. 194 (2): 257-75.
13. Kawabata, A. *et al.* (2011) Analysis of a Neutralizing Antibody for Human Herpesvirus 6B Reveals a Role for Glycoprotein Q1 in Viral Entry. J Virol. 85: 12962-71.
14. Cornfine, S. *et al.* (2011) The kinesin KIF9 and reggie/flotillin proteins regulate matrix degradation by macrophage podosomes. Mol Biol Cell. 22: 202-15.
15. Luo, S. *et al.* (2015) Contribution of N-linked glycans on HSV-2 gB to cell-cell fusion and viral entry. Virology. 483: 72-82.
16. El Kasmi, I. & Lippé, R. (2015) Herpes simplex virus 1 gN partners with gM to modulate the viral fusion machinery. J Virol. 89 (4): 2313-23.
17. Chia, R. *et al.* (2014) Phosphorylation of LRRK2 by casein kinase 1α regulates trans-Golgi clustering via differential interaction with ARHGEF7. Nat Commun. 5: 5827.
18. Vorobyeva, A.G. *et al.* (2014) Cyclopamine modulates γ-secretase-mediated cleavage of amyloid precursor protein by altering its subcellular trafficking and lysosomal degradation. J Biol Chem. 289 (48): 33258-74.
19. Wang Z *et al.* (2014) A newly identified myomegalin isoform functions in Golgi microtubule organization and ER-Golgi transport. J Cell Sci. 127 (22): 4904-17.
20. Ikawa Y *et al.* (2015) *In vitro* functional correction of Hermansky-Pudlak Syndrome type-1 by lentiviral-mediated gene transfer. Mol Genet Metab. 114 (1): 62-5.
21. Ioannou, M.S. *et al.* (2015) DENND2B activates Rab13 at the leading edge of migrating cells and promotes metastatic behavior. J Cell Biol. 208 (5): 629-48.
22. Gottschalk, E.Y. & Meneses, P.I. (2015) A Dual Role for the Nonreceptor Tyrosine Kinase Pyk2 during the Intracellular Trafficking of Human Papillomavirus 16. J Virol. 89 (17): 9103-14.
23. Crevenna, A.H. *et al.* (2016) Secretory cargo sorting by Ca2+-dependent Cab45 oligomerization at the trans-Golgi network. J Cell Biol. 213 (3): 305-14.
24. Matrone, C. *et al.* (2016) Mannose 6-Phosphate Receptor Is Reduced in -Synuclein Overexpressing Models of Parkinsons Disease. PLoS One. 11 (8): e0160501.
25. Ketteler, R. *et al.* (2017) Image-based siRNA screen to identify kinases regulating Weibel-Palade body size control using electroporation. Sci Data. 4: 170022.
26. Lukhele, S. & Cohen É.A. (2017) Conserved residues within the HIV-1 Vpu transmembrane-proximal hinge region modulate BST2 binding and antagonism. Retrovirology. 14 (1): 18.
27. Haugsten, E.M. *et al.* (2016) Proximity Labeling Reveals Molecular Determinants of

FGFR4 Endosomal Transport. J Proteome Res. 15 (10): 3841-55.
28. Paquin, N. *et al.* (2016) The Conserved VPS-50 Protein Functions in Dense-Core Vesicle Maturation and Acidification and Controls Animal Behavior. Curr Biol. 26 (7): 862-71.
29. Sugden, S.M. *et al.* (2017) HIV-1 Vpu Downmodulates ICAM-1 Expression, Resulting in Decreased Killing of Infected $CD4^+$ T Cells by NK Cells. J Virol. 91 (8): pii: e02442-16.
30. DiGiuseppe, S. *et al.* (2015) Topography of the Human Papillomavirus Minor Capsid Protein L2 during Vesicular Trafficking of Infectious Entry. J Virol. 89 (20): 10442-52.
31. Cabukusta, B. *et al.* (2017) Ceramide phosphoethanolamine synthase SMSr is a target of Caspase-6 during apoptotic cell death. Biosci Rep. Jun 28. pii: BSR20170867 [Epub ahead of print]
32. Luchsinger, C. *et al.* (2018) Functional disruption of the Golgi apparatus protein ARF1 sensitizes MDA-MB-231 breast cancer cells to the antitumor drugs Actinomycin D and Vinblastine through ERK and AKT signaling. PLoS One. 13 (4): e0195401.
33. Piccolo, P. *et al.* (2019) Geleophysic dysplasia: novel missense variants and insights into ADAMTSL2 intracellular trafficking. Mol Genet Metab Rep. 21: 100504.
34. Ayala, I. *et al.* (2019) GRASP65 controls Golgi position and structure during G2/M transition by regulating the stability of microtubules. Traffic. 20 (10): 785-802.
35. Cavieres, V.A. *et al.* (2020) Human Golgi phosphoprotein 3 is an effector of RAB1A and RAB1B. PLoS One. 15 (8): e0237514.

Further Reading	1. Ponnambalam, S. *et al.* (1996) Primate homologues of rat TGN38: primary structure, expression and functional implications. J Cell Sci. 109 (Pt 3): 675-85.
Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10048 available at: https://www.bio-rad-antibodies.com/SDS/AHP500 10048
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Sheep IgG (H/L) (5184-2304...) Biotin

North & South America Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com
Worldwide Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com
Europe Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M379026:210323'

Printed on 02 Sep 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint