

Datasheet: AHP500GT

BATCH NUMBER 174181

Description:	SHEEP ANTI HUMAN TGN46
Specificity:	TGN46
Other names:	TGOLN2
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	10 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen (1)	▪			0.1ug/ml - 1ug/ml
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting	▪			0.1ug/ml - 1ug/ml
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) Fixation with 3% paraformaldehyde or methanol/acetone is recommended.

Target Species	Human
Species Cross Reactivity	Reacts with: Primate N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
Product Form	Purified IgG - liquid

Antiserum Preparation Antisera to human TGN46 were raised by repeated immunisation of sheep with highly purified antigen. Purified IgG prepared by affinity chromatography.

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	<0.1% Sodium Azide (NaN ₃) 0.5% Bovine Serum Albumin 25% Glycerol
Approx. Protein Concentrations	IgG concentration 0.25 mg/ml
Immunogen	Recombinant human TGN46.
External Database Links	UniProt: O43493 Related reagents Entrez Gene: 10618 TGN46 Related reagents
Synonyms	TGN46, TGN51
RRID	AB_2203291
Specificity	Sheep anti Human TGN46 antibody recognizes trans-Golgi network integral membrane protein 2 (TGN46), also known as TGN38 homolog, TGN46, TGN48 or trans-Golgi network protein TGN51. TGN46 is a 437 amino acid glycoprotein localized to the trans-Golgi network. TGN46 has been reported as being the best available marker for human trans-Golgi network. TGN46 is a heavily glycosylated protein of around 110-120 kDa. Multiple isoforms of TGN46 are generated by alternative splicing differing in sequence at the C-terminal portion. Sheep anti Human TGN46 antibody is expected to recognize all identified isoforms.
References	1. Prescott AR <i>et al.</i> (1997) Distinct compartmentalization of TGN46 and beta 1,4-galactosyltransferase in HeLa cells. Eur J Cell Biol. 72 (3): 238-46. 2. van Dam, E.M. <i>et al.</i> (2002) Dynamin-dependent transferrin receptor recycling by endosome-derived clathrin-coated vesicles. Mol Biol Cell. 13: 169-82. 3. Salahpour, A. <i>et al.</i> (2004) Homodimerization of the beta2-adrenergic receptor as a prerequisite for cell surface targeting. J Biol Chem. 279 (32): 33390-7. 4. Drakesmith, H. <i>et al.</i> (2005) HIV-1 Nef down-regulates the hemochromatosis protein HFE, manipulating cellular iron homeostasis. Proc Natl Acad Sci U S A. 102 (31): 11017-22. 5. Mills, I.G. <i>et al.</i> (2005) Huntingtin interacting protein 1 modulates the transcriptional activity of nuclear hormone receptors. J Cell Biol. 170 (2): 191-200. 6. Mills, I.G. <i>et al.</i> (2005) Huntingtin interacting protein 1 modulates the transcriptional activity of nuclear hormone receptors. J Cell Biol. 170: 191-200.

7. Vuillier, F. *et al.* (2005) Lower levels of surface B-cell-receptor expression in chronic lymphocytic leukemia are associated with glycosylation and folding defects of the mu and CD79a chains. [Blood. 105 \(7\): 2933-40.](#)
8. Berarducci, B. *et al.* (2006) Essential functions of the unique N-terminal region of the varicella-zoster virus glycoprotein E ectodomain in viral replication and in the pathogenesis of skin infection. [J Virol. 80: 9481-96.](#)
9. Edwards, T.L. *et al.* (2009) Endogenous spartin (SPG20) is recruited to endosomes and lipid droplets and interacts with the ubiquitin E3 ligases AIP4 and AIP5. [Biochem J. 423 \(1\): 31-9.](#)
10. Esk, C. *et al.* (2010) The clathrin heavy chain isoform CHC22 functions in a novel endosomal sorting step. [J Cell Biol. 188: 131-44.](#)
11. Vleck, S.E. *et al.* (2010) Anti-glycoprotein H antibody impairs the pathogenicity of varicella-zoster virus in skin xenografts in the SCID mouse model. [J Virol. 84: 141-52.](#)
12. Sadaoka, T. *et al.* (2010) Characterization of the varicella-zoster virus ORF50 gene, which encodes glycoprotein M. [J Virol. 84: 3488-502.](#)
13. Hauser, H. *et al.* (2010) HIV-1 Vpu and HIV-2 Env counteract BST-2/tetherin by sequestration in a perinuclear compartment. [Retrovirology. 7: 51.](#)
14. Roberts, R.C. *et al.* (2010) Mistargeting of SH3TC2 away from the recycling endosome causes Charcot-Marie-Tooth disease type 4C. [Hum Mol Genet. 19: 1009-18.](#)
15. Fairn, G.D. *et al.* (2011) High-resolution mapping reveals topologically distinct cellular pools of phosphatidylserine. [J Cell Biol. 194 \(2\): 257-75.](#)
16. Vleck, S.E. *et al.* (2011) Structure-function analysis of varicella-zoster virus glycoprotein H identifies domain-specific roles for fusion and skin tropism. [Proc Natl Acad Sci U S A. 108 \(45\): 18412-7.](#)
17. Oliver, S.L. *et al.* (2011) Mutagenesis of varicella-zoster virus glycoprotein I (gI) identifies a cysteine residue critical for gE/gI heterodimer formation, gI structure, and virulence in skin cells. [J Virol. 85 \(9\): 4095-110.](#)
18. Petit, S.J. *et al.* (2011) Analysis of the human immunodeficiency virus type 1 M group Vpu domains involved in antagonizing tetherin. [J Gen Virol. 92 \(Pt 12\): 2937-48.](#)
19. Zuckerman, D.M. *et al.* (2011) Differential regulation of two palmitoylation sites in the cytoplasmic tail of the beta1-adrenergic receptor. [J Biol Chem. 286: 19014-23.](#)
20. Laufman, O. *et al.* (2011) The COG complex interacts directly with Syntaxin 6 and positively regulates endosome-to-TGN retrograde transport. [J Cell Biol. 194 \(3\): 459-72.](#)
21. Uchida, Y. *et al.* (2011) Intracellular phosphatidylserine is essential for retrograde membrane traffic through endosomes. [Proc Natl Acad Sci U S A. 108 \(38\): 15846-51.](#)
22. Cheng, S.B. *et al.* (2011) Down-modulation of the G-protein-coupled Estrogen Receptor, GPER, from the Cell Surface Occurs via a trans-Golgi-Proteasome Pathway. [J Biol Chem. 286: 22441-55.](#)
23. Kwon, S. and Christian, J.L. (2011) Sortilin Associates with Transforming Growth Factor- β Family Proteins to Enhance Lysosome-mediated Degradation. [J Biol Chem. 286: 21876-85.](#)
24. Kawabata, A. *et al.* (2011) Analysis of a Neutralizing Antibody for Human Herpesvirus 6B Reveals a Role for Glycoprotein Q1 in Viral Entry. [J Virol. 85: 12962-71.](#)
25. Cornfine, S. *et al.* (2011) The kinesin KIF9 and reggie/flotillin proteins regulate matrix degradation by macrophage podosomes. [Mol Biol Cell. 22: 202-15.](#)
26. Vorobyeva, A.G. *et al.* (2014) Cyclopamine modulates γ -secretase-mediated cleavage of amyloid precursor protein by altering its subcellular trafficking and lysosomal

- degradation. [J Biol Chem. 289 \(48\): 33258-74.](#)
27. Chia, R. *et al.* (2014) Phosphorylation of LRRK2 by casein kinase 1 α regulates trans-Golgi clustering via differential interaction with ARHGEF7. [Nat Commun. 5: 5827.](#)
 28. Wang Z *et al.* (2014) A newly identified myomegalin isoform functions in Golgi microtubule organization and ER-Golgi transport. [J Cell Sci. 127 \(22\): 4904-17.](#)
 29. DiGiuseppe, S. *et al.* (2015) Topography of the Human Papillomavirus Minor Capsid Protein L2 during Vesicular Trafficking of Infectious Entry. [J Virol. 89 \(20\): 10442-52.](#)
 30. Gottschalk, E.Y. & Meneses, P.I. (2015) A Dual Role for the Nonreceptor Tyrosine Kinase Pyk2 during the Intracellular Trafficking of Human Papillomavirus 16. [J Virol. 89 \(17\): 9103-14.](#)
 31. Ioannou, M.S. *et al.* (2015) DENND2B activates Rab13 at the leading edge of migrating cells and promotes metastatic behavior. [J Cell Biol. 208 \(5\): 629-48.](#)
 32. El Kasmi, I. & Lippé, R. (2015) Herpes simplex virus 1 gN partners with gM to modulate the viral fusion machinery. [J Virol. 89 \(4\): 2313-23.](#)
 33. Luo, S. *et al.* (2015) Contribution of N-linked glycans on HSV-2 gB to cell-cell fusion and viral entry. [Virology. 483: 72-82.](#)
 34. Ikawa Y *et al.* (2015) *In vitro* functional correction of Hermansky-Pudlak Syndrome type-1 by lentiviral-mediated gene transfer. [Mol Genet Metab. 114 \(1\): 62-5.](#)
 35. Crevenna, A.H. *et al.* (2016) Secretory cargo sorting by Ca²⁺-dependent Cab45 oligomerization at the trans-Golgi network. [J Cell Biol. 213 \(3\): 305-14.](#)
 36. Matrone, C. *et al.* (2016) Mannose 6-Phosphate Receptor Is Reduced in α -Synuclein Overexpressing Models of Parkinsons Disease. [PLoS One. 11 \(8\): e0160501.](#)
 37. Haugsten, E.M. *et al.* (2016) Proximity Labeling Reveals Molecular Determinants of FGFR4 Endosomal Transport. [J Proteome Res. 15 \(10\): 3841-55.](#)
 38. Paquin, N. *et al.* (2016) The Conserved VPS-50 Protein Functions in Dense-Core Vesicle Maturation and Acidification and Controls Animal Behavior. [Curr Biol. 26 \(7\): 862-71.](#)
 39. Ketteler, R. *et al.* (2017) Image-based siRNA screen to identify kinases regulating Weibel-Palade body size control using electroporation. [Sci Data. 4: 170022.](#)
 40. Lukhele, S. & Cohen É.A. (2017) Conserved residues within the HIV-1 Vpu transmembrane-proximal hinge region modulate BST2 binding and antagonism. [Retrovirology. 14 \(1\): 18.](#)
 41. Sugden, S.M. *et al.* (2017) HIV-1 Vpu Downmodulates ICAM-1 Expression, Resulting in Decreased Killing of Infected CD4⁺ T Cells by NK Cells. [J Virol. 91 \(8\): pii: e02442-16.](#)
 42. Cabukusta, B. *et al.* (2017) Ceramide phosphoethanolamine synthase SMSr is a target of Caspase-6 during apoptotic cell death. [Biosci Rep. 37 \(4\): BSR20170867.](#)
 43. Luchsinger, C. *et al.* (2018) Functional disruption of the Golgi apparatus protein ARF1 sensitizes MDA-MB-231 breast cancer cells to the antitumor drugs Actinomycin D and Vinblastine through ERK and AKT signaling. [PLoS One. 13 \(4\): e0195401.](#)
 44. Ayala, I. *et al.* (2019) GRASP65 controls Golgi position and structure during G2/M transition by regulating the stability of microtubules. [Traffic. 20 \(10\): 785-802.](#)
 45. Piccolo, P. *et al.* (2019) Geleophysic dysplasia: novel missense variants and insights into ADAMTSL2 intracellular trafficking. [Mol Genet Metab Rep. 21: 100504.](#)
 46. Cavieres, V.A. *et al.* (2020) Human Golgi phosphoprotein 3 is an effector of RAB1A and RAB1B. [PLoS One. 15 \(8\): e0237514.](#)
 47. Sakuma, C. *et al.* (2021) Identification of SYS1 as a Host Factor Required for Shiga Toxin-Mediated Cytotoxicity in Vero Cells [Int J Mol Sci 22 \(9\): 4936.](#)

48. Besemer, A.S. *et al.* (2021) Receptor-mediated endocytosis 8 (RME-8)/DNAJC13 is a novel positive modulator of autophagy and stabilizes cellular protein homeostasis. [Cell Mol Life Sci. 78 \(2\): 645-60.](#)
49. Stoneham, C.A. *et al.* (2021) A combined EM and proteomic analysis places HIV-1 Vpu at the crossroads of retromer and ESCRT complexes: PTPN23 is a Vpu-cofactor. [PLoS Pathog. 17 \(11\): e1009409.](#)
50. Yoshimura, A. *et al.* (2021) Branched Actin Maintains Acetylated Microtubule Network in the Early Secretory Pathway [Cells. 11 \(1\): 15.](#)
51. Bracci, N. *et al.* (2022) Rift Valley fever virus Gn V5-epitope tagged virus enables identification of UBR4 as a Gn interacting protein that facilitates Rift Valley fever virus production. [Virology. 567: 65-76.](#)
52. Zhang, J. *et al.* (2022) SARS-CoV-2 triggers Golgi fragmentation via down-regulation of GRASP55 to facilitate viral trafficking [Preprint] [bioRxiv 09 Mar \[Epub ahead of print\].](#)
53. Hertel, A. *et al.* (2022) USP32-regulated LAMTOR1 ubiquitination impacts mTORC1 activation and autophagy induction. [Cell Rep. 41 \(10\): 111653.](#)
54. Park, D. *et al.* (2023) Synaptic vesicle proteins and ATG9A self-organize in distinct vesicle phases within synapsin condensates. [Nat Commun. 14 \(1\): 455.](#)
55. Wang, J. *et al.* (2023) The clathrin adaptor complex-1 and Rab12 regulate post-Golgi trafficking of wild-type epidermal growth factor receptor (EGFR). [J Biol Chem. : 102979.](#)
56. Nishino, M. *et al.* (2023) Histone methyltransferase SUV39H1 regulates the Golgi complex via the nuclear envelope-spanning LINC complex. [PLoS One. 18 \(7\): e0283490.](#)
57. Hao, H. *et al.* (2020) Golgi-associated microtubules are fast cargo tracks and required for persistent cell migration. [EMBO Rep. 21 \(3\): e48385.](#)
58. Fujimoto, T. *et al.* (2018) Parkinson's disease-associated mutant LRRK2 phosphorylates Rab7L1 and modifies trans-Golgi morphology. [Biochem Biophys Res Commun. 495 \(2\): 1708-15.](#)
59. Venditti, R. *et al.* (2019) Molecular determinants of ER-Golgi contacts identified through a new FRET-FLIM system. [J Cell Biol. 218 \(3\): 1055-65.](#)
60. Lopes-da-Silva, M. *et al.* (2019) A GBF1-Dependent Mechanism for Environmentally Responsive Regulation of ER-Golgi Transport. [Dev Cell. 49 \(5\): 786-801.e6.](#)
61. Wakana, Y. *et al.* (2021) The ER cholesterol sensor SCAP promotes CARTS biogenesis at ER-Golgi membrane contact sites. [J Cell Biol. 220 \(1\): e202002150.](#)
62. Hieda, M. *et al.* (2021) The SUN2-nesprin-2 LINC complex and KIF20A function in the Golgi dispersal. [Sci Rep. 11 \(1\): 5358.](#)
63. Barbosa, N.S. *et al.* (2018) ESCRT machinery components are required for Orthobunyavirus particle production in Golgi compartments. [PLoS Pathog. 14 \(5\): e1007047.](#)
64. Afram, E. *et al.* (2023) The η -secretase-derived APP fragment η CTF is localized in Golgi, endosomes and extracellular vesicles and contributes to A β production. [Cell Mol Life Sci. 80 \(4\): 97.](#)
65. van der Beek, J. *et al.* (2024) Loss of the HOPS complex disrupts early-to-late endosome transition, impairs endosomal recycling and induces accumulation of amphisomes. [Mol Biol Cell. : mbcE23080328 \[Epub ahead of print\]](#)
66. Wirth, M. *et al.* (2021) Phosphorylation of the LIR Domain of SCOC Modulates ATG8 Binding Affinity and Specificity. [J Mol Biol. 433 \(13\): 166987.](#)
67. Hecht, T.K. *et al.* (2020) Fam20C regulates protein secretion by Cab45 phosphorylation. [J Cell Biol. 219 \(6\): e201910089.](#)

68. Jermusyk, A. *et al.* (2021) A 584 bp deletion in CTRB2 inhibits chymotrypsin B2 activity and secretion and confers risk of pancreatic cancer. [Am J Hum Genet. 108 \(10\): 1852-65.](#)
69. Lujan, P. *et al.* (2024) Sorting of secretory proteins at the trans-Golgi network by human TGN46. [Elife. 12: RP91708.](#)
70. Concha, J.O. *et al.* (2024) Rab27a GTPase and its effector Myosin Va are host factors required for efficient Oropouche virus cell egress. [PLoS Pathog. 20 \(8\): e1012504.](#)
71. Stein, A. *et al.* (2025) The Parkinson Disease-Associated Mutant DNAJC13(N855S) Leads to Its Accelerated Degradation and Negatively Affects Macroautophagy and Retromer Complex-Mediated Dynamics. [J Cell Physiol. 240 \(7\): e70074.](#)
72. Pirozzi, M. *et al.* (2025) Cargoes move from cis to trans-Golgi compartments and concentrate in the TGN before exiting. [EMBO Rep. 26 \(19\): 4742-65.](#)

Further Reading	1. Ponnambalam, S. <i>et al.</i> (1996) Primate homologues of rat TGN38: primary structure, expression and functional implications. J Cell Sci. 109 (Pt 3): 675-85.
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10048 available at: https://www.bio-rad-antibodies.com/SDS/AHP500GT
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Sheep IgG (H/L) (5184-2304...) [Biotin](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M437984:250320'

Printed on 21 Aug 2026