

Datasheet: AHP727 BATCH NUMBER 154276

| RABBIT ANTI PUMA |
|---------------------|
| PUMA |
| PUMA |
| Purified |
| Polyclonal Antibody |
| Polyclonal IgG |
| 0.1 mg |
| |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|--------------------------------|-----|----|----------------|--------------------|
| Flow Cytometry | | | • | |
| Immunohistology - Frozen | | | • | |
| Immunohistology - Paraffin (1) | • | | | 10 ug/ml |
| ELISA | | | • | |
| Immunoprecipitation | | | • | |
| Western Blotting | • | | | 2ug/ml - 4ug/ml |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

| Target Species | Human | |
|-----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------|
| Species Cross Reactivity | Based on sequence similarity, is expected to react with: Mouse N.B. Antibody reactivity and working conditions may vary betwee reactivity is derived from testing within our laboratories, peer-repersonal communications from the originators. Please refer to further information. | veen species. Cross eviewed publications or |
| Product Form | Purified IgG - liquid | |

| Anticorum rioparation | rabbits with highly purified antigen. Purified IgG was prepared chromatography. | • |
|--------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------|
| Buffer Solution | Phosphate buffered saline | |
| Preservative Stabilisers | 0.02% Sodium Azide | |
| Approx. Protein Concentrations | IgG concentration 1.0 mg/ml | |
| Immunogen | Synthetic peptide ARARQEGSSPEPVEG corresponding to the binding component 3. | e N-terminus of human Bcl-2 |
| External Database Links | UniProt: Q9BXH1 Related reagents Entrez Gene: 27113 BBC3 Related reagents | |
| Synonyms | PUMA | |
| RRID | AB_324241 | |
| Specificity | Rabbit anti PUMA antibody recognizes the p53 - upregulated also known as Bcl-2 binding component 3, JFY-1 or Bcl-2. PU kDa pro-apoptotic member of the BCL-2 family. | • • |
| | Bcl-2 binding component 3 is localized to the mitochondria, bin cytochrome C release and apoptosis in response to p53 (Folliof PUMA have been described due to alternative splicing. Rate expected to recognize the canonical isoform 1, also known as expected to recognize any of the other isoforms of PUMA. | s <i>et al.</i> 2013). Four isoforms obit anti PUMA antibody is |
| Histology Positive Control Tissue | Human breast carcinoma | |
| Western Blotting | AHP727 detects a band of approximately 23kDa in K562 cell I | ysates. |
| References | 1. Naderi, E.H. <i>et al.</i> (2009) Activation of cAMP signaling inhib apoptosis in BCP-ALL cells through abrogation of p53 accumb 608-18. | <u>-</u> |
| Further Reading | 1. Nakano, K. & Vousden, K.H. (2001) PUMA, a novel proapo p53. Mol Cell. 7 (3): 683-94. | ptotic gene, is induced by |
| Storage | Store at +4°C or at -20°C if preferred. | |

Antiserum Preparation Antisera to human Bcl-2 binding component 3 were raised by repeated immunisations of

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

| Guarantee | 12 months from date of despatch |
|----------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Health And Safety Information | Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/AHP727 10040 |
| Regulatory | For research purposes only |

Related Products

Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...) **FITC**

Goat Anti Rabbit IgG (Fc) (STAR121...) Biotin, FITC, HRP

Sheep Anti Rabbit IgG (STAR35...)

Goat Anti Rabbit IgG (H/L) (STAR124...) HRP

Recommended Useful Reagents

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A) TidyBlot WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

North & South Tel: +1 800 265 7376 Worldwide Fax: +1 919 878 3751 America

Email: antibody_sales_us@bio-rad.com

Tel: +44 (0)1865 852 700 Europe Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M364288:200529'

Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint