

Datasheet: AHP727

Description:	RABBIT ANTI PUMA
Specificity:	PUMA
Other names:	JFY-1
Format:	Purified
Product Type:	Polyclonal Antibody
lsotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .					
		Yes	No	Not Determined	Suggested Dilution	
	Flow Cytometry			•		
	Immunohistology - Frozen			•		
	Immunohistology - Paraffin (1)	•			10 ug/ml	
	ELISA					
	Immunoprecipitation					
	Western Blotting				2ug/ml - 4ug/ml	
	necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls. (1)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections.Sodium citrate buffer pH 6.0 is recommended for this purpose.					
Target Species	Human					
Species Cross Reactivity	Based on sequence similarity, is expected to react with:Mouse N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.					
Product Form	Purified IgG - liquid					

Antiserum Preparation Antisera to human Bcl-2 binding component 3 were raised by repeated immunisations of

	rabbits with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.02% Sodium Azide		
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml		
Immunogen	Synthetic peptide ARARQEGSSPEPVEG corresponding to the N-terminus of human Bcl-2 binding component 3.		
External Database Links	UniProt: Q9BXH1 Related reagents Entrez Gene: 27113 BBC3 Related reagents		
Synonyms	PUMA		
RRID	AB_324241		
Specificity	 Rabbit anti PUMA antibody recognizes the p53 - upregulated modulator of apoptosis, also known as Bcl-2 binding component 3, JFY-1 or Bcl-2. PUMA is a 193 amino acid ~20 kDa pro-apoptotic member of the BCL-2 family. Bcl-2 binding component 3 is localized to the mitochondria, binds BCL-2 and induces cytochrome C release and apoptosis in response to p53 (Follis <i>et al.</i> 2013). Four isoforms of PUMA have been described due to alternative splicing. Rabbit anti PUMA antibody is expected to recognize the canonical isoform 1, also known as PUMA alpha, it is not expected to recognize any of the other isoforms of PUMA. 		
Specificity Histology Positive Control Tissue	 also known as Bcl-2 binding component 3, JFY-1 or Bcl-2. PUMA is a 193 amino acid ~20 kDa pro-apoptotic member of the BCL-2 family. Bcl-2 binding component 3 is localized to the mitochondria, binds BCL-2 and induces cytochrome C release and apoptosis in response to p53 (Follis <i>et al.</i> 2013). Four isoforms of PUMA have been described due to alternative splicing. Rabbit anti PUMA antibody is expected to recognize the canonical isoform 1, also known as PUMA alpha, it is not 		
Histology Positive	 also known as Bcl-2 binding component 3, JFY-1 or Bcl-2. PUMA is a 193 amino acid ~20 kDa pro-apoptotic member of the BCL-2 family. Bcl-2 binding component 3 is localized to the mitochondria, binds BCL-2 and induces cytochrome C release and apoptosis in response to p53 (Follis <i>et al.</i> 2013). Four isoforms of PUMA have been described due to alternative splicing. Rabbit anti PUMA antibody is expected to recognize the canonical isoform 1, also known as PUMA alpha, it is not expected to recognize any of the other isoforms of PUMA. 		
Histology Positive Control Tissue	 also known as Bcl-2 binding component 3, JFY-1 or Bcl-2. PUMA is a 193 amino acid ~20 kDa pro-apoptotic member of the BCL-2 family. Bcl-2 binding component 3 is localized to the mitochondria, binds BCL-2 and induces cytochrome C release and apoptosis in response to p53 (Follis <i>et al.</i> 2013). Four isoforms of PUMA have been described due to alternative splicing. Rabbit anti PUMA antibody is expected to recognize the canonical isoform 1, also known as PUMA alpha, it is not expected to recognize any of the other isoforms of PUMA. Human breast carcinoma 		
Histology Positive Control Tissue Western Blotting	 also known as Bcl-2 binding component 3, JFY-1 or Bcl-2. PUMA is a 193 amino acid ~20 kDa pro-apoptotic member of the BCL-2 family. Bcl-2 binding component 3 is localized to the mitochondria, binds BCL-2 and induces cytochrome C release and apoptosis in response to p53 (Follis <i>et al.</i> 2013). Four isoforms of PUMA have been described due to alternative splicing. Rabbit anti PUMA antibody is expected to recognize the canonical isoform 1, also known as PUMA alpha, it is not expected to recognize any of the other isoforms of PUMA. Human breast carcinoma AHP727 detects a band of approximately 23kDa in K562 cell lysates. 1. Naderi, E.H. <i>et al.</i> (2009) Activation of cAMP signaling inhibits DNA damage-induced apoptosis in BCP-ALL cells through abrogation of p53 accumulation. Blood. 114 (3): 		

	short term use (up to 4 weeks) and store the remaining aliquots at -20°C.				
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.				
Guarantee	12 months from date of despatch				
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/AHP727 10040				
Regulatory	For research purposes only				

Related Products

Recommended Secondary Antibodies

 Sheep Anti Rabbit IgG (STAR34...)
 FITC

 Sheep Anti Rabbit IgG (STAR35...)
 RPE

 Goat Anti Rabbit IgG (Fc) (STAR121...)
 Biotin, FITC, HRP

 Sheep Anti Rabbit IgG (STAR36...)
 DyLight®488, DyLight®680, DyLight®800

 Goat Anti Rabbit IgG (H/L) (STAR124...)
 HRP

Recommended Useful Reagents

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A) TidyBlot WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	id.com	Email: antibody_sales_uk@bio-r	ad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M382642:210513'

Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint