

# Datasheet: AHP767 BATCH NUMBER 151215

Description:	RABBIT ANTI HUMAN EGF
Specificity:	EGF
Format:	Purified
<b>Product Type:</b>	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

## **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="https://www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			•	
Immunohistology - Frozen			•	
Immunohistology - Paraffin (1)	•			5.0ug/ml
ELISA	•			0.5 - 2.0ug/ml
Immunoprecipitation				
Western Blotting	•			0.1 - 0.2ug/ml
Functional Assays	•			<0.1ug/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1)This product requires heat-mediated pre-treatment of paraffin sections prior to staining. Citrate buffer pH 6.0 is recommended for this purpose

Target Species	Human
Product Form	Purified IgG - lyophilised
Reconstitution	Reconstitute with 0.1ml distilled water.
	For long term storage the addition of 0.09% sodium azide is recommended.
	N.B. For functional studies do not add azide.
	Care should be taken during reconstitution as the protein may appear as a film at the
	bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution.

Antiserum Preparation	Antisera to human EGF were raised by repeated immunisations of rabbits with highly purified antigen.  Purified IgG prepared by affinity chromatography.
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	None present.
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0mg/ml after reconstitution
Immunogen	Recombinant human EGF ( <u>PHP030A</u> ).
External Database Links	UniProt: P01133 Related reagents
	Entrez Gene:  1950 EGF Related reagents
RRID	AB_2095966
Specificity	
оресписку	<b>Rabbit anti Human EGF</b> recognizes human Epidermal Growth Factor, a potent stimulato and regulator of the proliferation of epidermal and epithelial cells, expressed by a variety of cells.
эреспісіту	and regulator of the proliferation of epidermal and epithelial cells, expressed by a variety
ELISA	and regulator of the proliferation of epidermal and epithelial cells, expressed by a variety of cells.  The effects of EGF are initiated through tyrosine kinase activity, following the binding of EGF to the extracellular domain of the ~170 kDa EGF receptor (EGFR), which also binds other EGF-like ligands with high affinity, including TGF-a (transforming growth factor alpha), VGF (vaccinia virus growth factor) and Betacellulin, expressed in the pancreas, small intestine and by certain tumour cells (Seno et al. 1996). Receptor activation through EGF binding triggers several signal transduction pathways, including the JAK/STAT and
	and regulator of the proliferation of epidermal and epithelial cells, expressed by a variety of cells.  The effects of EGF are initiated through tyrosine kinase activity, following the binding of EGF to the extracellular domain of the ~170 kDa EGF receptor (EGFR), which also binds other EGF-like ligands with high affinity, including TGF-a (transforming growth factor alpha), VGF (vaccinia virus growth factor) and Betacellulin, expressed in the pancreas, small intestine and by certain tumour cells (Seno et al. 1996). Receptor activation through EGF binding triggers several signal transduction pathways, including the JAK/STAT and P13K/AKT pathways.  This purified human EGF antibody may be used in an indirect ELISA or as the capture reagent in a sandwich ELISA with a biotinylated human EGF antibody (AHP767B) as the
ELISA Histology Positive	and regulator of the proliferation of epidermal and epithelial cells, expressed by a variety of cells.  The effects of EGF are initiated through tyrosine kinase activity, following the binding of EGF to the extracellular domain of the ~170 kDa EGF receptor (EGFR), which also binds other EGF-like ligands with high affinity, including TGF-a (transforming growth factor alpha), VGF (vaccinia virus growth factor) and Betacellulin, expressed in the pancreas, small intestine and by certain tumour cells (Seno et al. 1996). Receptor activation through EGF binding triggers several signal transduction pathways, including the JAK/STAT and P13K/AKT pathways.  This purified human EGF antibody may be used in an indirect ELISA or as the capture reagent in a sandwich ELISA with a biotinylated human EGF antibody (AHP767B) as the detection reagent and recombinant human EGF (PHP030A) as the standard.

### 7709-12.

Storage	Prior to reconstitution store at -20°C.
	After reconstitution store at -20°C.
	This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody.
	Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety	Material Safety Datasheet documentation #10162 available at:
Information	https://www.bio-rad-antibodies.com/SDS/AHP767
	10162
Regulatory	For research purposes only

# Related Products

# **Recommended Secondary Antibodies**

Goat Anti Rabbit IgG (Fc) (STAR121...) Biotin, FITC, HRP Sheep Anti Rabbit IgG (STAR35...) Goat Anti Rabbit IgG (H/L) (STAR124...) HRP

## **Recommended Useful Reagents**

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A) TidyBlot WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

Worldwide

America Fax: +1 919 878 3751

North & South Tel: +1 800 265 7376

Email: antibody\_sales\_us@bio-rad.com

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M372537:200706'

Email: antibody\_sales\_uk@bio-rad.com

#### Printed on 29 Feb 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint