

Datasheet: AHP781

BATCH NUMBER 164580

Description:	RABBIT ANTI HUMAN INTERLEUKIN-8
Specificity:	IL-8
Other names:	CXCL8
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin (1)	▪			0.25 - 1.0ug/ml
ELISA	▪			0.5 - 2.0ug/ml
Immunoprecipitation			▪	
Western Blotting	▪			0.1 - 0.2ug/ml
Immunofluorescence	▪			
Functional Assays	▪			2.0 - 3.0ug/ml

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own systems using appropriate negative/positive controls.

(1) This product requires enzyme mediated antigen retrieval prior to staining of parrafin sections. Proteinase K is recomended for this purpose.

Target Species	Human
Product Form	Purified IgG - lyophilized
Reconstitution	Reconstitute with 0.1ml distilled water Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution.

For long term storage the addition of 0.09% sodium azide is recommended.
N.B. For functional studies do not add sodium azide

Antiserum Preparation	Antisera to human IL-8 were raised by repeated immunisations of rabbits with highly purified antigen. Purified IgG prepared by affinity chromatography.
------------------------------	---

Buffer Solution	Phosphate buffered saline
------------------------	---------------------------

Preservative Stabilisers	None present
---------------------------------	--------------

Carrier Free	Yes
---------------------	-----

Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
---------------------------------------	-----------------------------

Immunogen	Recombinant human IL-8 (PHP055)
------------------	---

External Database Links	UniProt: P10145 Related reagents Entrez Gene: 3576 IL8 Related reagents
--------------------------------	--

Synonyms	CXCL8
-----------------	-------

RRID	AB_2264854
-------------	------------

Specificity	<p>Rabbit anti Human Interleukin-8 antibody recognizes human IL-8 (Interleukin-8), a 72 amino acid pro-inflammatory CXC chemokine designated CXCL8, secreted by endothelial cells and monocytes, which acts as a major chemoattractant for neutrophils and also T cells and basophils.</p> <p>IL-8, first isolated from osteosarcoma cells, contains the ELR-motif (N-Terminal Glu-Leu-Arg amino acid sequence) and signals through the CXCR1 and CXCR2 receptors. Previous nomenclature for IL-8 includes neutrophil activating protein 1(NAP-1), granulocyte chemotactic protein 1 (GCP-1), monocyte-derived neutrophil-activating peptide (MONAP) and protein 3-10C.</p>
--------------------	--

ELISA	This purified human IL-8 antibody may be used in an indirect ELISA or as the capture reagent in a sandwich ELISA with a biotinylated human IL-8 antibody (AHP781B) as the detection reagent and recombinant human IL-8 (PHP055) as the standard.
--------------	--

Western Blotting	This purified human IL-8 antibody may used in Western Blotting applications under either reducing or non-reducing conditions with recombinant human IL-8 (PHP055) as the positive control.
-------------------------	--

References	1. Vazquez, G.R. <i>et al.</i> (2018) Adipocytes Promote Early Steps of Breast Cancer Cell
-------------------	--

Dissemination via Interleukin-8. [Front Immunol. 9: 1767.](#)

2. Fu, X. *et al.* (2023) High FOXA1 levels induce ER transcriptional reprogramming, a pro-metastatic secretome, and metastasis in endocrine-resistant breast cancer. [Cell Rep. 42 \(8\): 112821.](#)

Further Reading	1. Mukaida, N. <i>et al.</i> (1989) Genomic structure of the human monocyte-derived neutrophil chemotactic factor IL-8. J Immunol. 143 (4): 1366-71.
------------------------	--

Storage	Prior to reconstitution store at -20°C. After reconstitution store at -20°C. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
----------------	--

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10294 available at: https://www.bio-rad-antibodies.com/SDS/AHP781 10294
--------------------------------------	---

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)

Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)

Sheep Anti Rabbit IgG (STAR35...) [RPE](#)

Recommended Useful Reagents

[ANTIGEN RETRIEVAL BUFFER, pH8.0 \(BUF025A\)](#)

[TidyBlot WESTERN BLOT DETECTION REAGENT:HRP \(STAR209P\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M402209:220718'

Printed on 15 Jul 2024