

## Datasheet: AHP781B

**BATCH NUMBER 100914**

<b>Description:</b>	RABBIT ANTI HUMAN INTERLEUKIN-8:Biotin
<b>Specificity:</b>	IL-8
<b>Other names:</b>	CXCL8
<b>Format:</b>	Biotin
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	50 µg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			0.25 - 1.0ug/ml
Immunoprecipitation			▪	
Western Blotting	▪			0.1 - 0.2ug/ml

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own systems using appropriate negative/positive controls.

<b>Target Species</b>	Human
<b>Product Form</b>	Purified IgG conjugated to Biotin - lyophilised
<b>Reconstitution</b>	Reconstitute with 0.5 ml sterile PBS containing 0.1% Bovine Serum Albumin. Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution. For long term storage the addition of 0.09% sodium azide is recommended.

**Antiserum Preparation** Antisera to human IL-8 were raised by repeated immunisations of rabbits with highly purified antigen. Purified IgG prepared by affinity chromatography.

<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	None present
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 0.1mg/ml after reconstitution
<b>Immunogen</b>	<a href="#">Recombinant human IL-8</a> (PHP055)
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P10145</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">3576</a>    IL8    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	CXCL8
<b>RRID</b>	AB_2126224
<b>Specificity</b>	<p><b>Rabbit anti Human Interleukin-8 antibody</b> recognizes human IL-8 (Interleukin-8), a 72 amino acid pro-inflammatory CXC chemokine designated CXCL8, secreted by endothelial cells and monocytes, which acts as a major chemoattractant for neutrophils and also T cells and basophils.</p> <p>IL-8, first isolated from osteosarcoma cells, contains the ELR-motif (N-Terminal Glu-Leu-Arg amino acid sequence) and signals through the CXCR1 and CXCR2 receptors. Cancer studies have demonstrated a role for IL-8 in the angiogenesis and growth of tumours. Previous nomenclature for IL-8 includes neutrophil activating protein 1(NAP-1), granulocyte chemotactic protein 1 (GCP-1), monocyte-derived neutrophil-activating peptide (MONAP) and protein 3-10C.</p>
<b>ELISA</b>	This biotinylated human IL-8 antibody may be used in a direct ELISA or as the detection reagent in a sandwich ELISA with a <a href="#">purified human IL-8 antibody</a> (AHP781) as the capture reagent and <a href="#">recombinant human IL-8</a> (PHP055) as the standard.
<b>Western Blotting</b>	This biotinylated human IL-8 antibody may be used in Western Blotting applications under either reducing or non-reducing conditions with <a href="#">recombinant human IL-8</a> (PHP055) as the positive control.
<b>References</b>	1. Rodriguez, G.V. <i>et al.</i> (2018) Adipocytes Promote Early Steps of Breast Cancer Cell Dissemination via Interleukin-8 <a href="#">Frontiers in Immunology. 9 [Epub ahead of print]</a> .
<b>Further Reading</b>	1. Mukaida, N. <i>et al.</i> (1989) Genomic structure of the human monocyte-derived neutrophil chemotactic factor IL-8. <a href="#">J Immunol. 143 (4): 1366-71.</a>

**Storage** Prior to reconstitution store at -20°C.  
After reconstitution store at -20°C.

This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10162 available at:  
<https://www.bio-rad-antibodies.com/SDS/AHP781B>  
10162

---

**Regulatory** For research purposes only

---

**North & South America** Tel: +1 800 265 7376  
Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700  
Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21  
Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M372550:200706'

**Printed on 18 Jan 2024**

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)