

Datasheet: AHP786

Description:	RABBIT ANTI HUMAN MDC
Specificity:	MDC
Other names:	CCL22
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			0.5ug/ml
Immunoprecipitation			▪	
Western Blotting	▪			0.1 - 0.2ug/ml
Functional Assays	▪			5.0 - 7.0ug/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - lyophilised
Reconstitution	Reconstitute with 0.1ml distilled water. For long term storage the addition of 0.09% sodium azide is recommended. NB: For functional studies do not add azide.
Antiserum Preparation	Antisera to human MDC were raised by repeated immunisations of rabbits with highly purified antigen. Purified IgG prepared by affinity chromatography.
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	None present
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0mg/ml

Immunogen	Recombinant human MDC.
External Database Links	<p>UniProt: O00626 Related reagents</p> <p>Entrez Gene: 6367 CCL22 Related reagents</p>
Synonyms	MDC, SCYA22
RRID	AB_808370
Specificity	<p>Rabbit anti Human MDC antibody recognizes human MDC, also known as macrophage-derived cytokine, CCL22, CC chemokine STCP-1, Small-inducible cytokine A22 or Stimulated T-cell chemotactic protein 1. MDC is a 93 amino acid, ~10.6 kDa member of the CC chemokine family expressed by macrophages, monocyte-derived dendritic cells, epithelial cells, fibroblasts and highly expressed in the thymus.</p> <p>MDC acts as a chemoattractant for monocytes, dendritic cells, activated T lymphocytes and NK cells, through binding to chemokine (C-C motif) receptor 4 (CCR4). CCR4 is selectively expressed by Th2 cells and several studies have revealed a link between MDC and another ligand of CCR4, known as activation-regulated chemokine (TARC/CCL17), with Th2-mediated inflammatory disorders such as atopic dermatitis and ulcerative colitis (Judge et al. 2001).</p>
Further Reading	1. Imai, T. <i>et al.</i> (1998) Macrophage-derived chemokine is a functional ligand for the CC chemokine receptor 4. J Biol Chem. 273 (3): 1764-8.
Storage	<p>Prior to reconstitution store at -20°C. After reconstitution store at -20°C.</p> <p>This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10162 available at: 10162: https://www.bio-rad-antibodies.com/uploads/MSDS/10162.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

- Sheep Anti Rabbit IgG (STAR34...) [FITC](#)
- Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)
- Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)
- Sheep Anti Rabbit IgG (STAR35...) [RPE](#)
- Sheep Anti Rabbit IgG (STAR36...) [DyLight@488](#), [DyLight@680](#), [DyLight@800](#)

Recommended Useful Reagents

TidyBlot WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M372556:200706'

Printed on 11 Aug 2020

© 2020 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)