

## Datasheet: AHP792

<b>Description:</b>	RABBIT ANTI HUMAN PDGF AA HOMODIMER
<b>Specificity:</b>	PDGF AA HOMODIMER
<b>Other names:</b>	PLATELET DERIVED GROWTH FACTOR AA
<b>Format:</b>	Purified
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin (1)	▪			
ELISA	▪			0.5ug/ml
Immunoprecipitation			▪	
Western Blotting	▪			0.1 - 0.2ug/ml
Functional Assays	▪			0.17 - 0.19ug/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

**(1) This product requires enzyme mediated antigen retrieval prior to staining of paraffin sections. Proteinase K is recommended for this purpose.**

<b>Target Species</b>	Human
<b>Product Form</b>	Purified IgG - lyophilised
<b>Reconstitution</b>	Reconstitute with 0.1ml distilled water. For long term storage the addition of 0.09% sodium azide is recommended. NB. For functional studies do not add azide.
<b>Antiserum Preparation</b>	Antisera to human PDGF-AA were raised by repeated immunisations of rabbits with highly purified antigen. Purified IgG prepared by affinity chromatography.

<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	None present
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0mg/ml after reconstitution
<b>Immunogen</b>	Recombinant human PDGF-AA.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P04085</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">5154</a> PDGFA    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	PDGF1
<b>RRID</b>	AB_844648
<b>Specificity</b>	<p><b>Rabbit anti Human PDGF AA Homodimer antibody</b> recognizes human PDGF-AA (Platelet derived growth factor A chain), a disulphide-linked alpha-chain homodimer and member of the PDGF family, released from platelet alpha-granules following trauma/platelet activation.</p> <p>Naturally occurring active forms of PDGF exist as dimers of two alpha chains (AA), two beta chains (BB), or a combination of both chains (AB), all of which signal through the high-affinity PDGF-Ralpha receptor, and the latter two of which can also signal through the PDGF-Rbeta receptor. PDGFs are potent mitogens for a variety of cells including those of the connective tissue, bone and cartilage, blood and smooth muscle cells, and are involved in several biological processes including chemotaxis, hyperplasia and embryonic neurogenesis.</p> <p>Increased expression of PDGF-AA has been associated with a diverse range of conditions including progression of osteosarcomas, persistent Cytomegalovirus infection in kidney allografts and mediation of IL-13-stimulated lung fibroblast proliferation in asthma.</p>
<b>Further Reading</b>	<ol style="list-style-type: none"> <li>1. Rorsman, F. <i>et al.</i> (1988) Structural characterization of the human platelet-derived growth factor A-chain cDNA and gene: alternative exon usage predicts two different precursor proteins. <a href="#">Mol Cell Biol. 8 (2): 571-7.</a></li> <li>2. Ingram, J.L. <i>et al.</i> (2004) IL-13 and IL-1beta promote lung fibroblast growth through coordinated up-regulation of PDGF-AA and PDGF-Ralpha. <a href="#">FASEB J. 18 (10): 1132-4.</a></li> </ol>
<b>Storage</b>	<p>Prior to reconstitution store at -20°C.</p> <p>After reconstitution store at -20°C.</p>

This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee** 6 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10162 available at: 10162: <https://www.bio-rad-antibodies.com/uploads/MSDS/10162.pdf>

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...) [FITC](#)

Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)

Sheep Anti Rabbit IgG (STAR35...) [RPE](#)

Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)

Sheep Anti Rabbit IgG (STAR36...) [DyLight®488](#), [DyLight®680](#), [DyLight®800](#)

### Recommended Useful Reagents

[ANTIGEN RETRIEVAL BUFFER, pH8.0 \(BUF025A\)](#)

[100x ANTIGEN RETRIEVAL BUFFER, pH8.0 \(BUF025C\)](#)

[TidyBlot WESTERN BLOT DETECTION REAGENT:HRP \(STAR209P\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M372561:200706'

**Printed on 16 Apr 2021**

---

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)