

## Datasheet: AHP849

<b>Description:</b>	GOAT ANTI HUMAN DEFENSIN BETA 2
<b>Specificity:</b>	DEFENSIN BETA 2
<b>Other names:</b>	BD-2
<b>Format:</b>	Purified
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Paraffin	▪			
ELISA	▪			0.5 - 2.0ug/ml
Immunoprecipitation			▪	
Western Blotting	▪			0.1 - 0.2ug/ml
Immunofluorescence			▪	
Functional Assays	▪			5.0 - 8.0ug/ml

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Human
<b>Product Form</b>	Purified IgG - lyophilised
<b>Reconstitution</b>	<p>Reconstitute with 1.0ml distilled water</p> <p>Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution. For long term storage the addition of 0.09% sodium azide is recommended.</p> <p>N.B. For functional studies do not add sodium azide</p>
<b>Preparation</b>	Antisera to human Defensin beta 2 were raised by repeated immunisation of goats with highly purified antigen. Purified IgG was prepared by affinity chromatography.

<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	None present
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 0.1 mg/ml after reconstitution
<b>Immunogen</b>	<a href="#">Recombinant human Defensin beta-2</a> (PHP161).
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">O15263</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">1673</a>    DEFB4A    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	DEFB102, DEFB2, DEFB4
<b>RRID</b>	AB_566948
<b>Specificity</b>	<p><b>Goat anti Human Defensin beta 2 polyclonal antibody</b> recognizes human Defensin beta-2 (BD-2), a ~4.3kDa <a href="#">cationic antimicrobial peptide</a>, also known as Beta-defensin 4A or Skin-antimicrobial peptide 1 (SAP-1) expressed primarily by epithelial cells of the respiratory tract and skin, and an important component of the innate immune response against microbial infections.</p> <p>Expression of BD-2 is induced during inflammation in response to bacterial products and cytokines, and is initially expressed in a precursor form, which is cleaved to release the C-Terminal active portion of the protein which is secreted by the neutrophil and binds to <a href="#">bacterial membranes</a> causing their disruption.</p>
<b>ELISA</b>	This purified human BD-2 antibody may be used in an indirect ELISA or as the capture reagent in a sandwich ELISA with a <a href="#">biotinylated human BD-2 antibody</a> (AHP849B) as the detection reagent and <a href="#">recombinant human BD-2</a> (PHP161) as the standard
<b>Western Blotting</b>	This purified human BD-2 antibody may be used in Western Blotting applications under either reducing or non-reducing conditions with <a href="#">recombinant human BD-2</a> (PHP161) as the positive control.
<b>References</b>	1. Crack, L.R. <i>et al.</i> (2012) Human antimicrobial peptides LL-37 and human $\beta$ -defensin-2 reduce viral replication in keratinocytes infected with varicella zoster virus. <a href="#">Clin Exp Dermatol. 37 (5): 534-43.</a>
<b>Further Reading</b>	<p>1. Bals, R. <i>et al.</i> (1998) Human beta-defensin 2 is a salt-sensitive peptide antibiotic expressed in human lung. <a href="#">J Clin Invest. 102 (5): 874-80.</a></p> <p>2. Harder, J. <i>et al.</i> (2000) Mucoid <i>Pseudomonas aeruginosa</i>, TNF-alpha, and IL-1beta, but</p>

not IL-6, induce human beta-defensin-2 in respiratory epithelia. [Am J Respir Cell Mol Biol. 22 \(6\): 714-21.](#)

---

**Storage** Prior to reconstitution store at -20°C.  
After reconstitution store at -20°C.

This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee** 6 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10162 available at:  
10162: <https://www.bio-rad-antibodies.com/uploads/MSDS/10162.pdf>

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) [FITC](#), [HRP](#)

### Recommended Useful Reagents

[ANTIGEN RETRIEVAL BUFFER, pH8.0 \(BUF025A\)](#)

[100x ANTIGEN RETRIEVAL BUFFER, pH8.0 \(BUF025C\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)  
'M372568:200706'

**Printed on 23 Mar 2021**

---

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)