

# Datasheet: AHP946 BATCH NUMBER 150796

Description:	GOAT ANTI DOG IgE	
Specificity:	lgE	
Format:	Purified	
Product Type:	Polyclonal Antibody	
Isotype:	Polyclonal IgG	
Quantity:	0.5 mg	

#### **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			•	
Immunohistology - Frozen			•	
Immunohistology - Paraffin				
ELISA	•			1/100 - 1/10,000
Immunoprecipitation			•	
Western Blotting	•			1/1000 - 1/30,000

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Dog
Product Form	Purified IgG - liquid

**Antiserum Preparation** Antisera to canine IgE were raised by repeated immunisations of goats with highly purified antigen. Purified IgG prepared by affinity chromatography.

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml

Native canine IgE.	
AB_567015	
Goat anti Dog IgE antibody recognises canine IgE. The antiboreact specifically with canine IgE by immunoelectrophoresis are cross reactivity was detected to canine IgG1, IgG2, IgA and IgCross react with IgE from other species.	nd ELISA. Less than 0.01%
Goat anti Dog IgE antibody has been reported for use in immu canine sections.	nohistology of frozen
1. Krachudel, J. <i>et al.</i> (2013) Luteal insufficiency in bitches as a autoimmune response against progesterone? Theriogenology.  2. Elders, R.C. <i>et al.</i> (2014) Recombinant canine IgE Fc and a protein bind to neoplastic canine mast cells. Vet Immunol Immunol 3. Couto N <i>et al.</i> (2016) Identification of vaccine candidate antiposeudintermedius by whole proteome characterization and ser analyses. J Proteomics. 133: 113-24.  4. Elders, R.C. <i>et al.</i> (2014) Recombinant canine IgE Fc and a protein bind to neoplastic canine mast cells. Vet Immunol Immunol Immunolisminate extract used in the treatment of canine atopic dermatitis Immunopathol. Aug 4 [Epub ahead of print]  6. Bizikova, P. <i>et al.</i> (2014) Serum autoantibody profiles of IgA pemphigus foliaceus. Vet Dermatol. 25 (5): 471-e75.	79: 1278-83  In IgE Fc-TRAIL fusion unopathol. 159: 29-40.  gens of Staphylococcus ological proteomic  In IgE Fc-TRAIL fusion unopathol. 159 (1-2): 29-40.
Store at +4°C. DO NOT FREEZE.  This product should be stored undiluted. Should this product correcommend microcentrifugation before use.	ontain a precipitate we
12 months from date of despatch	
Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/AHP946">https://www.bio-rad-antibodies.com/SDS/AHP946</a> 10040	
For research purposes only	
	Goat anti Dog IgE antibody recognises canine IgE. The antibreact specifically with canine IgE by immunoelectrophoresis are cross reactivity was detected to canine IgG1, IgG2, IgA and Igcross react with IgE from other species.  Goat anti Dog IgE antibody has been reported for use in immuneration sections.  1. Krachudel, J. et al. (2013) Luteal insufficiency in bitches as a autoimmune response against progesterone? Theriogenology.  2. Elders, R.C. et al. (2014) Recombinant canine IgE Fc and a protein bind to neoplastic canine mast cells. Vet Immunol Immunol 3. Couto N et al. (2016) Identification of vaccine candidate antipseudintermedius by whole proteome characterization and ser analyses. J Proteomics. 133: 113-24.  4. Elders, R.C. et al. (2014) Recombinant canine IgE Fc and a protein bind to neoplastic canine mast cells. Vet Immunol Immunolish Moya, R.et al. (2016) Immunoproteomic characterization of farinae extract used in the treatment of canine atopic dermatitis Immunopathol. Aug 4 [Epub ahead of print]  6. Bizikova, P. et al. (2014) Serum autoantibody profiles of IgA pemphigus foliaceus. Vet Dermatol. 25 (5): 471-e75.  Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. Should this product or recommend microcentrifugation before use.  12 months from date of despatch  Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/AHP946 10040

## **Related Products**

## **Recommended Secondary Antibodies**

Rabbit Anti Goat IgG (Fc) (STAR122...) FITC, HRP

Email: antibody\_sales\_us@bio-rad.com

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_uk@bio-rad.com

Email: antibody\_sales\_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

### Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint