

Datasheet: ANNEX100F

Description:	ANNEXIN V:FITC ASSAY KIT
Name:	ANNEXIN V KIT
Format:	FITC
Product Type:	Kits
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525

Preservative	0.02% Sodium Azide (NaN ₃)
Stabilisers	1% Bovine Serum Albumin

External Database Links

UniProt:

[P08758](#) [Related reagents](#)

Entrez Gene:

[308](#) ANXA5 [Related reagents](#)

Synonyms	ANX5, ENX2, PP4
-----------------	-----------------

Product Information

This test employs the property of Annexin V to bind to the membrane phospholipid phosphatidylserine (PS) in the presence of Ca²⁺. PS is exposed at the cell surface during the early stages of apoptosis. Detection of PS is a very sensitive method for detecting cells entering apoptosis, at a time point considerably ahead of nuclear changes such as DNA degradation.

The conjugation protocol used to prepare this product has not changed the native phospholipid binding properties of Annexin V. This protocol is designed to measure apoptosis easily and quickly in a sample of suspended cells.

[View our complete list of formats and sizes of Annexin V kits](#)

Reagents In The Kit Annexin V:FITC 1 x 0.5ml vial
 Propidium Iodide 1 x 1.6ml vial at 20ug/ml
 Binding buffer 1 x 50ml vial at x 4 concentrate

-
- Instructions For Use**
- 1) Dilute the binding buffer 1:4 in distilled water (50ml binding buffer +150ml distilled water)
 - 2) Wash cells in PBS by gentle shaking or by pipetting up and down.
 - 3) Resuspend cells in 200ul pre-diluted binding buffer, adjusting to a cell density of 2-5 x 10⁵ cells/ml.
 - 4) Add 5ul Annexin V:FITC to 195ul of the cell suspension prepared in step 3.
 - 5) Mix and incubate for 10 minutes in the dark, at room temperature.
 - 6) Wash cells in 200ul of pre-diluted binding buffer.
 - 7) Resuspend cells in 190ul pre-diluted binding buffer.
 - 8) Add 10ul of the Propidium Iodide solution.
 - 9) Analyse by flow cytometry.

The flow cytometer is preferably set such that the Mean Fluorescence Intensity of the Annexin V negative population is between 1 and 10. Optimal parameter settings can be found using a positive control. For a positive control, incubate the cells with 3% formaldehyde in buffer during 30 minutes on ice. Wash away the formaldehyde and suspend the cells in cold binding buffer at 2-5 x 10⁵ cells/ml. Proceed with step 2 as described above.

References

1. Lu, K.H. *et al.* (2010) *In Vitro* and *In Vivo* Apoptosis-Inducing Antileukemic Effects of Mucuna macrocarpa Stem Extract on HL-60 Human Leukemia Cells. [Integr Cancer Ther. 9: 298-308.](#)
2. Yen, J.H. *et al.* (2010) Glycine tomentella Hayata inhibits IL-1 β and IL-6 production, inhibits MMP-9 activity, and enhances RAW264.7 macrophage clearance of apoptotic cells. [J Biomed Sci. 17: 83.](#)
3. Chen, C.W. *et al.* (2010) The signals of FGFs on the neurogenesis of embryonic stem cells. [J Biomed Sci.17:33.](#)
4. Smith, K. *et al.* (2011) Mono- and tri-cationic porphyrin-monoclonal antibody conjugates: photodynamic activity and mechanism of action. [Immunology. 132: 256-65.](#)

5. Lu, K.H. *et al.* (2012) Synergistic Apoptosis-Inducing Antileukemic Effects of Arsenic Trioxide and Mucuna macrocarpa Stem Extract in Human Leukemic Cells via a Reactive Oxygen Species-Dependent Mechanism. [Evid Based Complement Alternat Med. 2012:921430.](#)
6. Koutsogiannaki, S. *et al.* (2015) Effects of cadmium and 17 β -estradiol on *Mytilus galloprovincialis* redox status. Prooxidant-antioxidant balance (PAB) as a novel approach in biomonitoring of marine environments. [Mar Environ Res. 103: 80-8.](#)
7. Iulia, A.I. *et al.* (2015) Inhibition of tumor necrosis factor alpha using RNA interference in oral squamous cell carcinoma. [J BUON. 20 \(4\): 1107-14.](#)
8. Hounkong, K. *et al.* (2015) Mechanisms of 1-hydroxy-2-hydroxymethylanthraquinone from *Coptosapelta flavescens* as an anti-giardial activity. [Acta Trop. 146: 11-6.](#)

Storage

Store at +4°C. DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

Guaranteed until date of expiry. Please see product label.

Health And Safety Information

Material Safety Datasheet documentation #10229 #10230 #10181 available at:
 Recombinant Annexin V (10229): <https://www.bio-rad-antibodies.com/uploads/MSDS/10229.pdf>
 Binding Buffer (10230): <https://www.bio-rad-antibodies.com/uploads/MSDS/10230.pdf>
 Propidium Iodide (10181): <https://www.bio-rad-antibodies.com/uploads/MSDS/10181.pdf>

Regulatory

For research purposes only

Related Products

Recommended Useful Reagents

[ANNEXIN V:FITC ASSAY KIT \(ANNEX300F\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
 'M393926:220120'

Printed on 21 Mar 2022