

Datasheet: ANNEX200APC BATCH NUMBER 166948

Description:	ANNEXIN V:APC ASSAY KIT
Name:	ANNEXIN V KIT
Format:	APC
Product Type:	Kits
Quantity:	200 TESTS

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols. Yes No Not Determined Suggested Dilution Flow Cytometry • Neat Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.							
Max Ex/Em	Fluorophore	Excitation Max (nr	n) Emission Max (nm)					
	APC	650	661					
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin							
External Database Links	UniProt: <u>P08758</u> <u>Related reagents</u> Entrez Gene: <u>308</u> ANXA5 <u>Related reagents</u>							
Synonyms	ANX5, ENX2, PP4							
Product Information	phosphatidylserine (P the early stages of ap	S) in the presence optosis. Detection o	n V to bind to the memb of Ca ²⁺ . PS is exposed of PS is a very sensitive onsiderably ahead of nu	at the cell surface during method for detecting				

The conjugation protocol used to prepare this product has not changed the native phospholipid binding properties of Annexin V. This protocol is designed to measure apoptosis easily and quickly in a sample of suspended cells.

View our complete list of formats and sizes of Annexin V kits

Annexin V:APC 200 tests					
Propidium Iodide Staining Solution 2X 100 tests					
10X Binding Buffer 100 ml					
1) Dilute the10X binding buffer to 1X in distilled water (1 ml binding buffer + 9 ml distilled water).					
2) Wash cells once in PBS by gentle shaking or pipetting up and down. Then wash once in 1X binding buffer.					
3) Resuspend cells in 1X binding buffer, adjusting to a cell density of 1-5 x 10 ⁶ cells/ml.					
4) Add 5 μ I of APC conjugated Annexin V to 100 μ I of the cell suspension.					
5) Mix and incubate for 10 to15 minutes at room temperature.					
6) Wash cells in 1X binding buffer.					
7) Resuspend cells in 200 μl of 1X binding buffer.					
8) Add 5 µl of Propidium Iodide Staining Solution.					
9) Analyse by flow cytometry within 4 hours. Store at 2-8°C in the dark.					
Store at +4°C. DO NOT FREEZE.					
This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.					
Guaranteed until date of expiry. Please see product label.					
Material Safety Datasheet documentation #10229 #10230 #10181 available at:					
https://www.bio-rad-antibodies.com/SDS/ANNEX200APC					
Recombinant Annexin V (10229)					
Binding Buffer (10230)					
Propidium lodide (10181)					

Related Products

Recommended Useful Reagents

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@	bio-rad.com	Email: antibody_sales_uk@bio	-rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M371228:200608'

Printed on 29 Feb 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint