

## Datasheet: ANNEX200PE BATCH NUMBER 164772

Description:	ANNEXIN V:PE ASSAY KIT
Name:	ANNEXIN V KIT
Format:	RPE
Product Type:	Kits
Quantity:	200 TESTS

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .    Yes No Not Determined Suggested Dilution   Flow Cytometry • Neat   Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.					
Max Ex/Em	Fluorophore	Excitation Max	(nm) E	mission Max (nm)		
	RPE 488nm laser	496		578		
Preservative Stabilisers	0.09% Sodium Azide (NaN <sub>3</sub> ) 1% Bovine Serum Albumin					
External Database Links	UniProt: <u>P08758</u> <u>Related reagents</u> Entrez Gene: <u>308</u> ANXA5 <u>Related reagents</u>					
Synonyms	ANX5, ENX2, PP4					
Product Information	the early stages of ap	PS) in the presen	ce of Ca on of PS	<sup>2+</sup> . PS is exposed is a very sensitive	at the cell surface during	

The conjugation protocol used to prepare this product has not changed the native phospholipid binding properties of Annexin V. This protocol is designed to measure apoptosis easily and quickly in a sample of suspended cells.

View our complete list of formats and sizes of Annexin V kits

Reagents In The Kit	Annexin V:PE	200 tests				
	7-AAD Viability Staining S	Solution 2X 100 tests				
	10X Binding Buffer	100 ml				
Instructions For Use	1) Dilute the10X Binding Buffer to 1X in distilled water (1 ml Binding Buffer + 9 ml distille water).					
	2) Wash cells once in PBS by gentle shaking or pipetting up and down. Then wash once in 1X Binding Buffer.					
	3) Resuspend cells in 1X	Binding Buffer, adjusting to a cell density of 1-5 x 10 <sup>6</sup> cells/ml				
	4) Add 5 µl of PE conjuga	ted Annexin V to 100 $\mu$ l of the cell suspension.				
	5) Mix and incubate for 10	) to15 minutes at room temperature.				
	6) Wash cells in 1X Bindir	ng Buffer.				
	7) Resuspend cells in 200	) μl of 1X Binding Buffer.				
	8) Add 5 µl of 7-AAD Viab temperature.	ility Staining Solution and incubate for 5 minutes at room				
	9) Analyse by flow cytome	etry within 4 hours. Store at 2-8°C in the dark.				
Storage	Store at +4°C. DO NOT F	REEZE.				
	This product should be sto	pred undiluted. Should this product contain a precipitate we				
	recommend microcentrifug	gation before use.				
Guarantee	Guaranteed until date of e	expiry. Please see product label.				
Health And Safety	Material Safety Datasheet	t documentation #10229 #10230 #10582 available at:				
Information	https://www.bio-rad-antibo	dies.com/SDS/ANNEX200PE				
	Recombinant Annexin V (	10229)				
	Binding Buffer (10230)					
	Binding Buffer (10230) 7-AAD (10582)					

**Related Products** 

## **Recommended Useful Reagents**

ANNEXIN V:PE ASSAY KIT (ANNEX50PE) ANNEXIN V:APC ASSAY KIT (ANNEX50APC) ANNEXIN V:APC ASSAY KIT (ANNEX200APC)

North & South	Tel: +1 800 265 7376	Worldwide
America	Fax: +1 919 878 3751	
	Email: antibody_sales_us@bio-rad.com	

Tel: +44 (0)1865 852 700 **Europe** Fax: +44 (0)1865 852 739 Email: antibody\_sales\_uk@bio-rad.com Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody\_sales\_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M371229:200608'

## Printed on 29 Feb 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint