

# Datasheet: ANNEX50APC

**BATCH NUMBER 170163**

<b>Description:</b>	ANNEXIN V:APC ASSAY KIT
<b>Name:</b>	ANNEXIN V KIT
<b>Format:</b>	APC
<b>Product Type:</b>	Kits
<b>Quantity:</b>	50 TESTS

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	APC	650	661

<b>Preservative</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Stabilisers</b>	1% Bovine Serum Albumin

### External Database Links

#### UniProt:

[P08758](#) [Related reagents](#)

#### Entrez Gene:

[308](#) ANXA5 [Related reagents](#)

<b>Synonyms</b>	ANX5, ENX2, PP4
-----------------	-----------------

### Product Information

This test employs the property of Annexin V to bind to the membrane phospholipid phosphatidylserine (PS) in the presence of Ca<sup>2+</sup>. PS is exposed at the cell surface during the early stages of apoptosis. Detection of PS is a very sensitive method for detecting cells entering apoptosis, at a time point considerably ahead of nuclear changes such as DNA degradation.

The conjugation protocol used to prepare this product has not changed the native phospholipid binding properties of Annexin V. This protocol is designed to measure apoptosis easily and quickly in a sample of suspended cells.

[View our complete list of formats and sizes of Annexin V kits](#)

<b>Reagents In The Kit</b>	Annexin V:APC 50 tests Propidium Iodide 100 tests 10X Binding Buffer 30 ml
<b>Instructions For Use</b>	<ol style="list-style-type: none"><li>1) Dilute the 10X binding buffer to 1X in distilled water (1 ml binding buffer + 9 ml distilled water).</li><li>2) Wash cells once in PBS by gentle shaking or pipetting up and down. Then wash once in 1X binding buffer.</li><li>3) Resuspend cells in 1X binding buffer, adjusting to a cell density of <math>1-5 \times 10^6</math> cells/ml.</li><li>4) Add 5 µl of APC conjugated Annexin V to 100 µl of the cell suspension.</li><li>5) Mix and incubate for 10 to 15 minutes at room temperature.</li><li>6) Wash cells in 1X binding buffer.</li><li>7) Resuspend cells in 200 µl of 1X binding buffer.</li><li>8) Add 5 µl of Propidium Iodide staining solution.</li><li>9) Analyse by flow cytometry within 4 hours. Store at 2-8°C in the dark.</li></ol>
<b>Storage</b>	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.
<b>Guarantee</b>	Guaranteed until date of expiry. Please see product label.
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10229 #10230 #10181 available at: <a href="https://www.bio-rad-antibodies.com/SDS/ANNEX50APC">https://www.bio-rad-antibodies.com/SDS/ANNEX50APC</a>
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Useful Reagents

[ANNEXIN V:APC ASSAY KIT \(ANNEX200APC\)](#)

Product inquiries: [www.bio-rad-antibodies.com/technical-support](http://www.bio-rad-antibodies.com/technical-support)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M371227:200608'

**Printed on 09 Jul 2025**

---

© 2025 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)