

Datasheet: ANNEX50APC

BATCH NUMBER 171299

Description:	ANNEXIN V:APC ASSAY KIT
Name:	ANNEXIN V KIT
Format:	APC
Product Type:	Kits
Quantity:	50 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	APC	650	661

Preservative	0.09% Sodium Azide (NaN ₃)
Stabilisers	1% Bovine Serum Albumin

External Database Links

UniProt:

[P08758](#) [Related reagents](#)

Entrez Gene:

[308](#) ANXA5 [Related reagents](#)

Synonyms	ANX5, ENX2, PP4
-----------------	-----------------

Product Information	This test employs the property of Annexin V to bind to the membrane phospholipid phosphatidylserine (PS) in the presence of Ca ²⁺ . PS is exposed at the cell surface during the early stages of apoptosis. Detection of PS is a very sensitive method for detecting cells entering apoptosis, at a time point considerably ahead of nuclear changes such as DNA degradation.
----------------------------	--

The conjugation protocol used to prepare this product has not changed the native phospholipid binding properties of Annexin V. This protocol is designed to measure apoptosis easily and quickly in a sample of suspended cells.

[View our complete list of formats and sizes of Annexin V kits](#)

Reagents In The Kit	Annexin V:APC 50 tests Propidium Iodide 100 tests 10X Binding Buffer 30 ml
Instructions For Use	1) Dilute the 10X binding buffer to 1X in distilled water (1 ml binding buffer + 9 ml distilled water). 2) Wash cells once in PBS by gentle shaking or pipetting up and down. Then wash once in 1X binding buffer. 3) Resuspend cells in 1X binding buffer, adjusting to a cell density of $1-5 \times 10^6$ cells/ml. 4) Add 5 µl of APC conjugated Annexin V to 100 µl of the cell suspension. 5) Mix and incubate for 10 to 15 minutes at room temperature. 6) Wash cells in 1X binding buffer. 7) Resuspend cells in 200 µl of 1X binding buffer. 8) Add 5 µl of Propidium Iodide staining solution. 9) Analyse by flow cytometry within 4 hours. Store at 2-8°C in the dark.
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	Guaranteed until date of expiry. Please see product label.
Health And Safety Information	Material Safety Datasheet documentation #10229 #10230 #10181 available at: https://www.bio-rad-antibodies.com/SDS/ANNEX50APC
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

[ANNEXIN V:APC ASSAY KIT \(ANNEX200APC\)](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M371227:200608'

Printed on 09 Jul 2025

© 2025 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)