

## Datasheet: APO010B BATCH NUMBER 157032

| Description:  | AUTOPHAGY ASSAY, RED DETECTION KIT |
|---------------|------------------------------------|
| Name:         | AUTOPHAGY ASSAY KIT, RED           |
| Format:       | Kit                                |
| Product Type: | Kits                               |
| Quantity:     | 200 TESTS                          |

## **Product Details**

| Applications        | This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> . |   |  |   |  |  |
|---------------------|--|---|--|---|--|--|
|                     |  | Yes   | No   | Not Determined  | Suggested Dilution   |  |
|                     | Flow Cytometry   | •   |  |   | Refer to Instructions For<br>Use   |  |
| Max Ex/Em           | Where this product has not been tested for use in a particular tern   necessarily exclude its use in such procedures.   Fluorophore Excitation Max (nm)   Emission Max (nm)  |   |  |   |  |  |
|                     | Red Probe  | 590   |  | 620   |  |  |
| Product Information | development of autop<br>Autophagy is a conse<br>own components suc<br>role in maintaining ho   | hagy in living<br>rved lysoson<br>h as proteins<br>meostasis by | g cells.<br>nal recycli<br>, lipids an<br>/ preventi | ng process by which o<br>d carbohydrates. Auto<br>ng the accumulation c | nd monitoring of in vitro<br>cells break down their<br>ophagy plays a critical<br>of damaged organelles by<br>mponents (Mizushima et |  |

role in maintaining homeostasis by preventing the accumulation of damaged organelles by disassembling unnecessary or dysfunctional cells and cellular components (Mizushima et al 2011). Autophagy occurs at low levels in the cell under normal conditions and can be rapidly upregulated during times of starvation or stress. Such degradation activities serve to provide nutrients (amino acids, nucleotides, fatty acids, etc.) and energy during periods of elevated bioenergetic demands (Mizushima et al 2011, Levine et al 2008). Another function of autophagy is to assist with the detection and destruction of intracellular pathogens (viruses, bacteria and parasites) (Levine et al 2011). Dysregulation of autophagy has been associated with many disease states including cancer, infection and degenerative diseases (Levine et al 2008). Autophagy is a dynamic process typically divided into three stages. During stage one, cytoplasmic components targeted for degradation are sequestered within a double-membrane phagopore (also called the isolation membrane). This results in the formation of a double-membrane vesicle called

|                                  | the autophagosome. During stage two, the autophagosome f<br>form the autolysosome. Degradation of the autophagosomal<br>three ( <u>Mizushima et al 2011</u> , <u>Hundeshagen et al 2011</u> ).   | •  |  |  |  |  |
|----------------------------------|--|--|--|--|--|--|
| Test Principle                   | Autophagy Probe, Red is a cell-permeant aliphatic molecule that fluoresces brightly when inserted in the lipid membranes of autophagosomes and autolysosomes. Autophagy Probe, Red can be readily detected by flow cytometry with optimal excitation at 590 nm and peak emission at 620 nm (ZE5 Cell Analyzer settings, 561 nm laser and 615/24 or 640/20 filter). |  |  |  |  |  |
| Reagents In The Kit              | Autophagy Probe, Red, 4 vials - lyophilized<br>Fixative, 6 ml  |  |  |  |  |  |
| Instructions For Use             | Instructions for use can be found at <u>www.bio-rad-antibodies.com/uploads/IFU/APO010.pdf</u>  |  |  |  |  |  |
| Storage                          | MULTIPLE STORAGE CONDITIONS APPLY ON ARRIVAL.<br>each unopened component) according to the storage instruct<br>label.<br>Store the Autophagy Probe, Red at -20°C. Once reconstitute<br>Probe, Red stock should be stored at -20°C for 6 months, pro-<br>repeated freezing and thawing.   | tions on each component  |  |  |  |  |
| Guarantee                        | Guaranteed until date of expiry. Please see product label.   |  |  |  |  |  |
| Health And Safety<br>Information | Material Safety Datasheet documentation #20402 #10498 av<br>https://www.bio-rad-antibodies.com/SDS/APO010B<br>Autophagy Probe, Red (20402)<br>Fixative (10498)   | ailable at:  |  |  |  |  |
| Regulatory                       | For research purposes only   |  |  |  |  |  |
|                                  | 3 3751 Fax: +44 (0)1865 852 739   _sales_us@bio-rad.com Email: antibody_sales_uk@bio-rad.com   | Tel: +49 (0) 89 8090 95 21<br>Fax: +49 (0) 89 8090 95 50<br>Email: antibody_sales_de@bio-rad.com |  |  |  |  |
| to find a batch/lot speci        | fic datasheet for this product, please use our online search tool at: bi<br>'M360112:191030'   | o-rau-antipoures.com/datasneets  |  |  |  |  |

Printed on 26 Jun 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint