

Datasheet: BUF034C BATCH NUMBER 167695

Description:	ELISA SYNBLOCK
Name:	ELISA SYNBLOCK
Format:	Ready To Use
Product Type:	Accessory Reagent
Quantity:	1000 ml

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .					
	ELISA	Yes	Νο	Not Determined	Suggested Dilution	
	Where this product has no	ot been te	sted for us	e in a particular tech		
	necessarily exclude its use a guide only. It is recomme system using appropriate r	ended tha	t the user	titrates the product for	•	
Product Form	Ready to use - liquid					
Buffer Solution	Phosphate buffered saline					
Preservative Stabilisers	0.05% Sodium Azide					
Product Information	ELISA Synblock is an EL with buffers containing ani signals in ELISA assays w	mal prote	ins (e.g. B	SA) and reduce non-	specific background	
Intended Use	ELISA SynBlock is a novel protein-free blocking buffer suitable for use in all ELISA formats requiring maximum blocking strength. With detergent and synthetic blocking agents, the inert nature of this unique buffer enables maximum reduction of non-specif binding and interference associated particularly with sandwich ELISA assays.					
	Additional molecular stabil environment for coating ar temperature and stored or	ntigen or o	capture an	tibody. Plates can be	•	

	N.B. SynBlock is not suitable for use on Immunlon [®] 2 plates. Bio-Rad recommends the use of <u>BUF033A</u> for this purpose.
Instructions For Use	1. Coat ELISA plate with antibody or antigen as required.
	2. After incubation, remove the coating solution and wash the plate x2 with wash buffer. <u>BUF031A</u> can be used for this purpose.
	3. Add 300-400ul of BUF034C and incubate for 2-24 hours. Use a volume equal to or greater than the volume of coating solution.
	4. After removal of the blocking buffer continue with the assay or dry the plate for long-term storage at +4°C.
References	 Afrough, B. <i>et al.</i> (2007) Identification and elimination of false-positives in an ELISA-based system for qualitative assessment of glycoconjugate binding using a selection of plant lectins. <u>Biotechniques. 43</u> (4): 458, 460, 462 passim. Dalley, D. <i>et al.</i> (2008) Development and evaluation of a gamma-interferon assay for tuberculosis in badgers (<i>Meles meles</i>). <u>Tuberculosis (Edinb). 88: 235-43</u>. Ahmed, R.R. <i>et al.</i> (2010) BACE1 and BACE2 enzymatic activities in Alzheimer's disease. <u>J Neurochem. 112: 1045-53</u>. Chambers, M.A. <i>et al.</i> (2009) Performance of TB immunodiagnostic tests in Eurasian badgers (<i>Meles meles</i>) of different ages and the influence of duration of infection on serological sensitivity. <u>BMC Vet Res. 5: 42</u>. Thompson, R. <i>et al.</i> (2011) Optimization of the enzyme-linked lectin assay for enhanced glycoprotein and glycoconjugate analysis. <u>Anal Biochem. 413: 114-22</u>. Kuramitz, H. <i>et al.</i> (2012) Multiplexed assay for proteins based on sequestration electrochemistry using the protein binding electroactive magnetic microbeads. <u>Anal Sci. 28</u> (1): 77. Dwek, M.V. <i>et al.</i> (2010) A sensitive assay to measure biomarker glycosylation demonstrates increased fucosylation of prostate specific antigen (PSA) in patients with prostate cancer compared with benign prostatic hyperplasia. <u>Clin Chim Acta. 411 (23-24)</u>; 1935-9. Verhelst, R. <i>et al.</i> (2010) The effects of plant polyphenols on enterotoxigenic <i>Escherichia coli</i> adhesion and toxin binding Livestock Science. 133 (1-3): 101-3 Verhelst, R. <i>et al.</i> (2013) <i>E. coli</i> heat labile toxin (LT) inactivation by specific polyphenols is aggregation dependent. <u>Vet Microbiol. 163 (3-4): 319-24</u>. Greenwell P <i>et al.</i> (2013) Lex coli edit improves motor performance but does not affect β-amyloid levels in a mouse model of Alzheimer's disease. <u>Brain Res. 1505: 61-7</u>. Abdul, H.M. <i>et al.</i> (2013) Leptin regulates amyloid β

	15. Wilcock, D.M. <i>et al.</i> (2015) Down syndrome individuals with Alzheimer's disease have a distinct neuroinflammatory phenotype compared to sporadic Alzheimer's disease.		
	Neurobiol Aging. 36 (9): 2468-74.		
	16. Chinthamani S <i>et al.</i> (2017) Macrophage inducible C-type lectin (Mincle) recognizes		
	glycosylated surface (S)-layer of the periodontal pathogen <i>Tannerella forsythia</i> . <u>PLoS</u> One. 12 (3): e0173394.		
17. LeVine, H. 3rd. <i>et al.</i> (2017) Down syndrome: age-dependence of PiB bir			
	postmortem frontal cortex across the lifespan. <u>Neurobiol Aging. 54: 163-9.</u>		
Storage	Store at +4°C.		
	DO NOT FREEZE		
Guarantee	Guaranteed until date of expiry. Please see product label.		
Health And Safety	Material Safety Datasheet documentation #10380 available at:		
Information	https://www.bio-rad-antibodies.com/SDS/BUF034C 10380		
Regulatory	For research purposes only		
Related Produ	cts		
Recommended U	seful Reagents		

5x ELISA COATING BUFFER (BUF030A) 10x ELISA WASH BUFFER (BUF031A)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-ra	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rac	Europe d.com	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com	
To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets						

'M353960:190618'

Printed on 29 Feb 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint