

Datasheet: BUF041B

Description:	MOUSE SEROBLOCK FcR
Name:	MOUSE SEROBLOCK FCR
Other names:	CD16/CD32
Format:	Reagent
Product Type:	Accessory Reagent
Clone:	FCR4G8
Isotype:	IgG2b
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	PU5 1.8 IOE7 Balb/c mouse cell line.

Product Information **Mouse Seroblock FcR** is a rat antibody that recognises mouse CD16 and CD32, cell surface proteins also known as FcRγIII and FcR&gamma:II, respectively. The function of these proteins is to bind IgG molecules via their Fc regions as part of the adaptive immune response. CD16 and/or CD32 are expressed by a wide variety of cells, including monocytes, macrophages, B lymphocytes, granulocytes, NK cells, dendritic cells and some activated T lymphocytes. The expression of CD16/CD32 antigens can lead to non-specific binding of test monoclonal antibodies in staining procedures, resulting in “high background” staining on a wide range of cells. This non-specific

binding may be blocked by pre-incubation of target cells with BUF041A, resulting in clearer staining. For direct analysis of CD16/32 expression this antibody is also available conjugated directly to FITC ([MCA2305F](#)), AlexaFluor488 (MCA2305A488), AlexaFluor647 ([MCA2305A647](#)) and RPE ([MCA2305PE](#)).

Instructions For Use In order to reduce Fc-receptor mediated binding of test antibodies the following procedure is recommended:-

1) Incubate the cell suspension (1 x 10⁶ cells in 100ul) with 1ug of BUF041B (1ul of undiluted reagent, or 10ul of a 1/10 dilution) for 5-10 minutes.

2) Add test antibody according to manufacturers instructions – Do not wash BUF041B off the cells. BUF041B is suitable for use in conjunction with test antibodies from any manufacturer or with in-house antibodies.*

3) Proceed with staining as usual.

* Care is needed in the design of experiments utilizing unconjugated anti-mouse antibodies, to ensure that the secondary antibody being used does not cross-react with SeroBlock FcR. Bio-Rad supply a range of isotype specific anti-mouse immunoglobulin antibodies that may be useful for this purpose.

References

1. Ivanovska, N.D *et al.* (2008) Properdin Deficiency in Murine Models of Nonseptic Shock. [The Journal of Immunology. 180: 6962-6969.](#)
2. Chen, H-F *et al.* (2009) A reduced oxygen tension (5%) is not beneficial for maintaining human embryonic stem cells in the undifferentiated state with short splitting intervals. [Human Reproduction. 24: 71-80.](#)
3. Birjandi, S.Z. *et al.* (2011) Alterations in marginal zone macrophages and marginal zone B cells in old mice. [J Immunol. 186: 3441-51.](#)
4. Cousins, F.L. *et al.* (2016) Evidence for a dynamic role for mononuclear phagocytes during endometrial repair and remodelling. [Sci Rep. 6: 36748.](#)

Storage

Store at +4°C or at -20°C if preferred.
Storage in frost-free freezers is not recommended.
This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

18 months from date of despatch.

Health And Safety Information

Material Safety Datasheet documentation #10040 available at:
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory

For research purposes only

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

'M352671:190405'

Printed on 20 May 2019
