Datasheet: BUF049C BATCH NUMBER 153658

| HISPEC ASSAY DILUENT | |
|----------------------|--|
| HISPEC ASSAY DILUENT | |
| Ready To Use | |
| Accessory Reagent | |
| N/A | |
| 500 ml | |
| | |

Product Details

| Applications | This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> . | | | | | | | | |
|-----------------------------|---|-----|----|----------------|--------------------|--|--|--|--|
| | ······ | Yes | No | Not Determined | Suggested Dilution | | | | |
| | ELISA | - | | | | | | | |
| | Immunoassay | • | | | | | | | |
| | Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls. | | | | | | | | |
| Product Form | Ready to use assay diluent buffer | | | | | | | | |
| Preservative Stabilisers | 0.1% Proclin™ 300 | | | | | | | | |
| Intended Use | HISPEC assay diluent works to reduce cross reactivity, non-specific binding and matrix effects in immunoassays such as ELISA, EIA, Western blotting, immuno-PCR, protein arrays, multianalyte immunoassays and immunohistochemistry. The effect of HISPEC assay diluent is dependent on assay system and the antibodies used. HISPEC assay diluent is used instead of a sample buffer or antibody dilution buffer | | | | | | | | |
| | locking of surfaces we | | | | | | | | |
| | recommend one of the blocking solutions within the Bio-Rad range – see "Use reagents" below. | | | | | | | | |

| ELISA: use as a dilution buffer for the specimen and for the detection antibodies Western blotting: use as dilution buffer for primary and secondary antibodies Immunohistochemistry: use as dilution buffer for primary and secondary antibodies |
|---|
| Protein arrays: use as dilution buffer for specimen and for the detection antibodies |
| Instructions For Use The buffer should be mixed thoroughly immediately before use. Dilution of the samples: The standards and samples for ELISA and protein arrays can be diluted with HISPEC assay diluent at 1:2 or higher. Standards and samples should be treated identically. |
| Dilution of antibodies: Antibodies can be diluted with HISPEC assay diluent in a user-defined manner, depending on the recommendation of the antibody supplier. |
| Expected results: In some cases a reduction in assay signal intensity may be observed when using HISPEC assay diluent. This does not result in a reduction in assay sensitivity due to the accompanying reduction in non-specific binding. |
| Signal intensity may be increased in some assay systems by use of higher concentrations of antibodies, or by dilution of the HISPEC assay diluent in distilled water (1:2) prior to use. However, these adjustments may also result in some associated increase in non-specific binding. |
| References 1. Eriksson, O. et al. (2017) Pancreatic imaging using an antibody fragment targeting the zinc transporter type 8: a direct comparison with radio-iodinated Exendin-4. Acta Diabetol. Oct 24 [Epub ahead of print]. |
| StorageStore at +4°C or at -20°C if preferred.Storage in frost-free freezers is not recommended.This product should be stored undiluted. Avoid repeated freezing and thawing. |
| Guarantee Guaranteed until date of expiry. Please see product label. |
| Acknowledgements Proclin [™] 300 is a trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow. |
| Health And Safety Material Safety Datasheet documentation #10265 available at: Information <u>https://www.bio-rad-antibodies.com/SDS/BUF049C</u> 10265 |
| Regulatory For research purposes only |

Related Products

Recommended Useful Reagents

| North & South | Tel: +1 800 265 7376 | Worldwide | Tel: +44 (0)1865 852 700 | Europe | Tel: +49 (0) 89 8090 95 21 |
|---------------|---------------------------|--------------|------------------------------|-----------|--------------------------------------|
| America | Fax: +1 919 878 3751 | | Fax: +44 (0)1865 852 739 | | Fax: +49 (0) 89 8090 95 50 |
| | Email: antibody_sales_us@ |)bio-rad.com | Email: antibody_sales_uk@bio | o-rad.com | Email: antibody_sales_de@bio-rad.com |

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M350303:190307'

Printed on 29 Feb 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint