

Datasheet: BUF09

BATCH NUMBER 157136

Description:	LEUCOPERM
Name:	LEUCOPERM
Format:	Reagent
Product Type:	Accessory Reagent
Quantity:	50 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			

LEUCOPERM reagents are intended for fixing cells in suspension with Reagent A and then permeabilising the cells with Reagent B. The specific formulations reduce background staining and allow simultaneous addition of permeabilisation medium and fluorochrome labelled antibodies.

Product Form	Reagent A - Fixation medium Reagent B - Permeabilisation medium
Preservative Stabilisers	Formaldehyde in Reagent A

Product Information Flow cytometric analyses with monoclonal antibodies have been restricted primarily to cell surface molecules. Intracellular structures such as cytoplasmic or nuclear enzymes, oncoproteins, cytokines, immunoglobulins etc. were largely excluded from such assays.

Also excluded from flow cytometric assays were cytoplasmic localizations of well established membrane molecules such as CD3 and CD22.

LEUCOPERM reagents allow intracellular antigen analysis with the same ease as surface antigens. The only prerequisite is the availability of suitable antibody conjugates. Most commercially available monoclonal antibody conjugates can be used with LEUCOPERM reagents. Some determinants are sensitive, however, to the fixation step involved. This and the optimal fixation time may have to be determined experimentally for each antibody conjugate.

Instructions For Use For the detection of cell cycle antigens such as Ki-67, PCNA and BrdU, methanol modification is recommended - see [protocol #F5](#).

1. Prepare cells in the appropriate manner. Adjust cell suspension to a concentration of 1×10^7 cells/ml in PBS/BSA. Whole blood samples may also be used. Bio-Rad recommend the use of EDTA anti-coagulant in these circumstances, although satisfactory results may be obtained using heparin or acid-citrate dextrose.
2. Add 100ul of cell suspension to the appropriate number of test tubes.
If required, perform staining of cell surface antigens at this stage. Following staining for the recommended period, wash cells once in PBS/BSA and discard supernatant.
3. Add 100ul of Reagent A (fixation medium, stored at room temperature).
4. Incubate for 15 minutes at room temperature.
5. Add 3ml PBS/BSA and centrifuge for 5 minutes at 300 x g. Remove supernatant.
6. Resuspend cells in 100ul of Reagent B (Permeabilization Medium).
7. Immediately add recommended volume of the appropriate directly conjugated antibody. Vortex and incubate for 30 minutes at room temperature.
If using an unconjugated primary antibody, wash in 3ml of PBS/BSA (as per step 5) and then repeat step 7 using an appropriate secondary antibody. There is no requirement to add further Leucoperm.
8. Wash once in PBS/BSA. Remove supernatant and resuspend cells in sheath fluid for immediate analysis or resuspend cells in 0.25ml of 0.5% formaldehyde and store them at 2-8°C in the dark. Analyse fixed cells within 24 hours.

References

1. Chiu, W.C. *et al.* (2009) Effects of dietary fish oil supplementation on cellular adhesion molecule expression and tissue myeloperoxidase activity in hypercholesterolemic mice with sepsis. [J Nutr Biochem. 20: 254-60.](#)
2. Grundy, M. *et al.* (2010) The FLT3 internal tandem duplication mutation is a secondary target of the aurora B kinase inhibitor AZD1152-HQPA in acute myelogenous leukemia cells. [Mol Cancer Ther. 9: 661-72.](#)
3. Taylor, L. *et al.* (2010) The effect of acute hypoxia on heat shock protein 72 expression and oxidative stress in vivo. [Eur J Appl Physiol. 109 \(5\): 849-55.](#)
4. Myles, A. *et al.* (2011) Expression of Toll-like receptors 2 and 4 is increased in peripheral blood and synovial fluid monocytes of patients with enthesitis-related arthritis subtype of juvenile idiopathic arthritis. [Rheumatology \(Oxford\). 50: 481-8.](#)
5. Osorio, Y. *et al.* (2011) Identification of small molecule lead compounds for visceral leishmaniasis using a novel ex vivo splenic explant model system [PLoS Negl Trop Dis. 5:e962.](#)
6. Jiang, W.J. *et al.* (2017) Structure-activity relationship of the inhibitory effects of flavonoids on nitric oxide production in RAW264.7 cells. [Bioorg Med Chem. 25 \(2\):](#)

[779-788.](#)

7. Kliminski, V. *et al.* (2016) Popdc1/Bves Functions in the Preservation of Cardiomyocyte Viability While Affecting Rac1 Activity and Bnip3 Expression. [J Cell Biochem. Nov 25. \[Epub ahead of print\]](#)
8. Parry, D.A. *et al.* (2016) A homozygous STIM1 mutation impairs store-operated calcium entry and natural killer cell effector function without clinical immunodeficiency. [J Allergy Clin Immunol. 137 \(3\): 955-7.e8.](#)
9. Suradhat, S. *et al.* (2015) A novel DNA vaccine for reduction of PRRSV-induced negative immunomodulatory effects: A proof of concept. [Vaccine. 33 \(32\): 3997-4003.](#)
10. Dishon, S. *et al.* (2017) Inhibition of Myeloid Differentiation Factor 88 Reduces Human and Mouse T-Cell Interleukin-17 and IFN γ Production and Ameliorates Experimental Autoimmune Encephalomyelitis Induced in Mice. [Front Immunol. 8: 615.](#)
11. Nie, H. *et al.* (2017) Phenotypic switch in lung interstitial macrophage polarization in an ovalbumin-induced mouse model of asthma. [Exp Ther Med. 14 \(2\): 1284-92.](#)
12. Arrieta-Villegas, C. *et al.* (2020) Immunogenicity and Protection against *Mycobacterium caprae* Challenge in Goats Vaccinated with BCG and Revaccinated after One Year. [Vaccines \(Basel\). 8 \(4\): 751.](#)
13. Alhuthali, H.M. *et al.* (2020) The natural alkaloid Jerantinine B has activity in acute myeloid leukemia cells through a mechanism involving c-Jun. [BMC Cancer. 20 \(1\): 629.](#)
14. Bairey, O. *et al.* (2010) Arsenic-trioxide-induced apoptosis of chronic lymphocytic leukemia cells. [Int J Lab Hematol. 32 \(1 Pt 1\): e77-85.](#)
15. Martelli, P. *et al.* (2021) Immune B cell responsiveness to single-dose intradermal vaccination against *Mycoplasma hyopneumoniae*. [Res Vet Sci. 141: 66-75.](#)
16. Hatzidaki, E. *et al.* (2021) A Novel Method for Colorectal Cancer Screening Based on Circulating Tumor Cells and Machine Learning. [Entropy \(Basel\). 23 \(10\): 1248.](#)
17. Martelli, P. *et al.* (2021) Immune B cell responsiveness to single-dose intradermal vaccination against *Mycoplasma hyopneumoniae*. [Res Vet Sci. 141: 66-75.](#)
18. Sanchez-Pino, M.D. (2022) Detection of Circulating and Tissue Myeloid-Derived Suppressor Cells (MDSC) by Flow Cytometry. [Methods Mol Biol. 2422: 247-61.](#)
19. Jeong, E.M. *et al.* (2022) Targeting RUNX1 as a novel treatment modality for pulmonary arterial hypertension. [Cardiovasc Res. Jan 09 \[Epub ahead of print\].](#)
20. Franzoni, G. *et al.* (2022) Analyses of the Impact of Immunosuppressive Cytokines on Porcine Macrophage Responses and Susceptibility to Infection to African Swine Fever Viruses. [Pathogens. 11 \(2\): 166.](#)

Storage	LEUCOPERM Cell Permeabilisation reagents should be stored and used at room temperature. DO NOT FREEZE. Do not use reagents if a precipitate forms or discolouration occurs.
----------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10187 #10509 available at: https://www.bio-rad-antibodies.com/SDS/BUF09 Reagent A - Fixation medium (10187) Reagent B - Permeabilisation medium (10509)
--------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Regulatory	For research purposes only
-------------------	----------------------------

North & South Tel: +1 800 265 7376

Worldwide Tel: +44 (0)1865 852 700

Europe Tel: +49 (0) 89 8090 95 21

America Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M374483:201109'

Printed on 17 May 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)