

Datasheet: BUF09C

Description:	LEUCOPERM
Name:	LEUCOPERM
Format:	Reagent
Product Type:	Accessory Reagent
Quantity:	1000 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			

LEUCOPERM reagents are intended for fixing cells in suspension with Reagent A and then permeabilising the cells with Reagent B. The specific formulations reduce background staining and allow simultaneous addition of permeabilisation medium and fluorochrome labelled antibodies.

Product Form	Reagent A - Fixation medium Reagent B - Permeabilisation medium
---------------------	--

Preservative Stabilisers	Formaldehyde in Reagent A
---------------------------------	---------------------------

Product Information Flow cytometric analyses with monoclonal antibodies have been restricted primarily to cell surface molecules. Intracellular structures such as cytoplasmic or nuclear enzymes, oncoproteins, cytokines, immunoglobulins etc. were largely excluded from such studies.

Also excluded from flow cytometric studies were cytoplasmic localisations of well established membrane molecules such as CD3 and CD22.

LEUCOPERM reagents allow intracellular antigen analysis with the same ease as surface antigens. The only prerequisite is the availability of suitable antibody conjugates. Most commercially available monoclonal antibody conjugates can be used with LEUCOPERM reagents. Some determinants are sensitive, however, to the fixation step involved. This and the optimal fixation time may have to be determined experimentally for each antibody conjugate.

Instructions For Use For the detection of cell cycle antigens such as Ki-67, PCNA and BrdU, methanol modification is recommended - see [protocol #F5](#).

1. Prepare cells in the appropriate manner. Adjust cell suspension to a concentration of 1×10^7 cells/ml in PBS/BSA. Whole blood samples may also be used. Bio-Rad recommend the use of EDTA anti-coagulant in these circumstances, although satisfactory results may be obtained using heparin or acid-citrate dextrose.
2. Add 100ul of cell suspension to the appropriate number of test tubes.
If required, perform staining of cell surface antigens at this stage. Following staining for the recommended period, wash cells once in PBS/BSA and discard supernatant.
3. Add 100ul of Reagent A (fixation medium, stored at room temperature).
4. Incubate for 15 minutes at room temperature.
5. Add 3ml PBS/BSA and centrifuge for 5 minutes at 300 x g. Remove supernatant.
6. Resuspend cells in 100ul of Reagent B (Permeabilization Medium).
7. Immediately add recommended volume of the appropriate directly conjugated antibody. Vortex and incubate for 30 minutes at room temperature.
If using an unconjugated primary antibody, wash in 3ml of PBS/BSA (as per step 5) and then repeat step 7 using an appropriate secondary antibody. There is no requirement to add further Leucoperm.
8. Wash once in PBS/BSA. Remove supernatant and resuspend cells in sheath fluid for immediate analysis or resuspend cells in 0.25ml of 0.5% formaldehyde and store them at 2-8°C in the dark. Analyse fixed cells within 24 hours.

References

1. Chiu, W.C. *et al.* (2009) Effects of dietary fish oil supplementation on cellular adhesion molecule expression and tissue myeloperoxidase activity in hypercholesterolemic mice with sepsis. [J Nutr Biochem. 20: 254-60.](#)
2. Grundy, M. *et al.* (2010) The FLT3 internal tandem duplication mutation is a secondary target of the aurora B kinase inhibitor AZD1152-HQPA in acute myelogenous leukemia cells. [Mol Cancer Ther. 9: 661-72.](#)
3. Taylor, L. *et al.* (2010) The effect of acute hypoxia on heat shock protein 72 expression and oxidative stress in vivo. [Eur J Appl Physiol. 109 \(5\): 849-55.](#)
4. Myles, A. *et al.* (2011) Expression of Toll-like receptors 2 and 4 is increased in peripheral blood and synovial fluid monocytes of patients with enthesitis-related arthritis subtype of juvenile idiopathic arthritis. [Rheumatology \(Oxford\). 50: 481-8.](#)
5. Osorio, Y. *et al.* (2011) Identification of small molecule lead compounds for visceral leishmaniasis using a novel ex vivo splenic explant model system [PLoS Negl Trop Dis. 5:e962.](#)
6. Jiang, W.J. *et al.* (2017) Structure-activity relationship of the inhibitory effects of flavonoids on nitric oxide production in RAW264.7 cells. [Bioorg Med Chem. 25 \(2\):](#)

[779-788.](#)

7. Kliminski, V. *et al.* (2016) Popdc1/Bves Functions in the Preservation of Cardiomyocyte Viability While Affecting Rac1 Activity and Bnip3 Expression. [J Cell Biochem. Nov 25. \[Epub ahead of print\]](#)
8. Parry, D.A. *et al.* (2016) A homozygous STIM1 mutation impairs store-operated calcium entry and natural killer cell effector function without clinical immunodeficiency. [J Allergy Clin Immunol. 137 \(3\): 955-7.e8.](#)
9. Suradhat, S. *et al.* (2015) A novel DNA vaccine for reduction of PRRSV-induced negative immunomodulatory effects: A proof of concept. [Vaccine. 33 \(32\): 3997-4003.](#)
10. Dishon, S. *et al.* (2017) Inhibition of Myeloid Differentiation Factor 88 Reduces Human and Mouse T-Cell Interleukin-17 and IFN γ Production and Ameliorates Experimental Autoimmune Encephalomyelitis Induced in Mice. [Front Immunol. 8: 615.](#)
11. Nie, H. *et al.* (2017) Phenotypic switch in lung interstitial macrophage polarization in an ovalbumin-induced mouse model of asthma. [Exp Ther Med. 14 \(2\): 1284-92.](#)
12. Arrieta-Villegas, C. *et al.* (2020) Immunogenicity and Protection against *Mycobacterium caprae* Challenge in Goats Vaccinated with BCG and Revaccinated after One Year. [Vaccines \(Basel\). 8 \(4\): 751.](#)
13. Alhuthali, H.M. *et al.* (2020) The natural alkaloid Jerantinine B has activity in acute myeloid leukemia cells through a mechanism involving c-Jun. [BMC Cancer. 20 \(1\): 629.](#)

Storage LEUCOPERM Cell Permeabilisation reagents should be stored and used at room temperature. DO NOT FREEZE. Do not use reagents if a precipitate forms or discolouration occurs.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10187 #10509 available at:
Reagent A - Fixation medium (10187): <https://www.bio-rad-antibodies.com/uploads/MSDS/10187.pdf>
Reagent B - Permeabilisation medium (10509): <https://www.bio-rad-antibodies.com/uploads/MSDS/10509.pdf>

Regulatory For research purposes only

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

From March 15, 2021, we will no longer supply printed datasheets with our products.
Look out for updates on how to access your digital version at [bio-rad-antibodies.com](https://www.bio-rad-antibodies.com)

'M374485:201109'

Printed on 02 Mar 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)