

Datasheet: C06SB

BATCH NUMBER 155163

Description:	DONKEY SERUM
Name:	DONKEY SERUM
Format:	Serum
Product Type:	Serum
Quantity:	100 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen	•			
Immunohistology - Paraffin	•			
Immunoassay	•			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Product Form	Serum - liquid
Preservative Stabilisers	0.1% Sodium Azide

Product Information

Donkey Serum is a sterile filtered preparation of donkey serum prepared from donkey whole blood following clotting.

References

- 1. Mou, H. *et al.* (2013) Generation of Multipotent Foregut Stem Cells from Human Pluripotent Stem Cells Cell Stem Cell. 10: 385-97.
- 2. Barghorn, S. *et al.* (2005) Globular amyloid beta-peptide oligomer a homogenous and stable neuropathological protein in Alzheimer's disease. J Neurochem. 95: 834-47.
- 3. Rashid, S.T. *et al.* (2010) Modeling inherited metabolic disorders of the liver using human induced pluripotent stem cells. <u>J Clin Invest. 120: 3127-36.</u>
- 4. McBride, P.A. *et al.* (1988) Immunostaining of scrapie cerebral amyloid plaques with antisera raised to scrapie-associated fibrils (SAF). <u>Neuropathol Appl Neurobiol. 14:</u> 325-36.
- 5. Vallier, L. et al. (2009) Early cell fate decisions of human embryonic stem cells and

mouse epiblast stem cells are controlled by the same signalling pathways. PLoS One. 4:

- 6. Debertin, G. et al. (2015) Tyrosine hydroxylase positive perisomatic rings are formed around various amacrine cell types in the mammalian retina. J Neurochem. 134 (3):
- 7. Pradillo, J.M. et al. (2017) Reparative effects of interleukin-1 receptor antagonist in young and aged/co-morbid rodents after cerebral ischemia. Brain Behav Immun. 61:
- 8. Das, R.G. et al. (2021) Neuron-specific spinal cord translatomes reveal a neuropeptide code for mouse dorsal horn excitatory neurons. Sci Rep. 11 (1): 5232.

St	ora	a	e

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the product. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10496 available at: https://www.bio-rad-antibodies.com/SDS/C06SB 10496
Regulatory	For research purposes only

North & South Tel: +1 800 265 7376 America

Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700

Europe Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M373106:200824'

Printed on 13 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint

Email: antibody_sales_us@bio-rad.com