

Datasheet: C12CA

BATCH NUMBER 155694

Description:	BABY RABBIT COMPLEMENT
Name:	BABY RABBIT COMPLEMENT
Format:	Serum
Product Type:	Serum
Quantity:	2 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Functional Assays (1)	▪			
Immunoassay	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) This product is not sold as sterile but can be sterilized by filtration if necessary. It is preferable to dilute the complement to a final working concentration before filtration in order to minimize loss of volume.

Product Form	Baby rabbit serum - lyophilised
Reconstitution	Reconstitute with 2ml ice cold distilled water
Preservative Stabilisers	None present
Product Information	Baby rabbit complement serum preparation is intended for use as a source of rabbit complement for cytotoxicity studies.
Instructions For Use	Use within one hour of reconstitution, keeping on ice throughout.

References

1. Council of Europe. (1983) Essential aspects of tissue typing: Preparation, use, storage of reagents and standardisation of complement. European Health Committee, Strasbourg, ISBN 92-871.224-4.
2. Edited by J.G.Ray. (1979) NIAID Manual of tissue typing techniques. Publication No.

NIH-80-545.

3. Anderson, L.D. Jr *et al.* (1999) Enhancement of graft-versus-tumor activity and graft-versus-host disease by pretransplant immunization of allogeneic bone marrow donors with a recipient-derived tumor cell vaccine. [Cancer Res. 59 \(7\): 1525-30.](#)
4. Mason, J.C. *et al.* (2002) Statin-induced expression of decay-accelerating factor protects vascular endothelium against complement-mediated injury. [Circ Res. 91 \(8\): 696-703.](#)
5. Lidington, E.A. *et al.* (2000) Induction of decay-accelerating factor by thrombin through a protease-activated receptor 1 and protein kinase C-dependent pathway protects vascular endothelial cells from complement-mediated injury. [Blood. 96 \(8\): 2784-92.](#)
6. De clerq, L. *et al.* (1997) An anti-adipocyte monoclonal antibody is cytotoxic to porcine preadipocytes *in vitro* and depresses the development of pig adipose tissue. [J Anim Sci. 75 \(7\): 1791-7.](#)
7. Hung, M.C. *et al.* (2011) The *Neisseria meningitidis* Macrophage Infectivity Potentiator Protein Induces Cross-Strain Serum Bactericidal Activity and Is a Potential Serogroup B Vaccine Candidate. [Infect Immun. 79: 3784-91.](#)
8. Lee, S.J. *et al.* (2012) Identification of a common immune signature in murine and human systemic Salmonellosis. [Proc Natl Acad Sci U S A. 109 \(13\): 4998-5003.](#)
9. Goh, Y.S. & MacLennan, C.A. (2013) Invasive African nontyphoidal *Salmonella* requires high levels of complement for cell-free antibody-dependent killing. [J Immunol Methods. 387 \(1-2\): 121-9.](#)
10. Li, S.H. *et al.* (2004) C-reactive protein upregulates complement-inhibitory factors in endothelial cells. [Circulation. 109: 833-6.](#)
11. Hyams, C. *et al.* (2010) *Streptococcus pneumoniae* resistance to complement-mediated immunity is dependent on the capsular serotype. [Infect Immun. 78: 716-25.](#)
12. Newcombe, J. *et al.* (2004) Infection with an avirulent *phoP* mutant of *Neisseria meningitidis* confers broad cross-reactive immunity. [Infect Immun. 72: 338-44.](#)
13. Mason, J.C. *et al.* (2002) bFGF and VEGF synergistically enhance endothelial cytoprotection via decay-accelerating factor induction. [Am J Physiol Cell Physiol. 282: C578-87.](#)
14. Sancho, D. *et al.* (2006) CD69 targeting differentially affects the course of collagen-induced arthritis. [J Leukoc Biol. 80: 1233-41.](#)
15. Hung MC *et al.* (2013) The adhesin complex protein (ACP) of *Neisseria meningitidis* is a new adhesin with vaccine potential. [MBio. 4 \(2\): pii: e00041-13.](#)
16. Goh YS *et al.* (2016) Bactericidal Immunity to *Salmonella* in Africans and Mechanisms Causing Its Failure in HIV Infection. [PLoS Negl Trop Dis. 10 \(4\): e0004604.](#)
17. Sawant S *et al.* (2016) Establishment of 3D Co-Culture Models from Different Stages of Human Tongue Tumorigenesis: Utility in Understanding Neoplastic Progression. [PLoS One. 11 \(8\): e0160615.](#)
18. Humbert MV *et al.* (2016) Vaccine Potential and Diversity of the Putative Cell Binding Factor (CBF, NMB0345/NEIS1825) Protein of *Neisseria meningitidis*. [PLoS One. 11 \(8\): e0160403.](#)
19. Nganje, C.N. *et al.* (2019) PepN is a non-essential, cell wall-localized protein that contributes to neutrophil elastase-mediated killing of *Streptococcus pneumoniae*. [PLoS One. 14 \(2\): e0211632.](#)
20. Dierckx de Casterlé I *et al.* (2018) Reduction of myeloid-derived suppressor cells reinforces the anti-solid tumor effect of recipient leukocyte infusion in murine

neuroblastoma-bearing allogeneic bone marrow chimeras. [Cancer Immunol Immunother. 67 \(4\): 589-603.](#)

21. Cuesta-Mateos, C. *et al.* (2020) CCR7 as a novel therapeutic target in t-cell PROLYMPHOCYTIC leukemia [Biomarker Research.8, 54.](#)

22. Valton, J. *et al.* (2018) A Versatile Safeguard for Chimeric Antigen Receptor T-Cell Immunotherapies. [Sci Rep. 8 \(1\): 8972.](#)

Storage

Prior to reconstitution store at +4°C. Following reconstitution store at +4°C for 1 hour or aliquot and store at -70°C for longer.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the product. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

Guaranteed until date of expiry. Please see product label.

Health And Safety Information

Material Safety Datasheet documentation #10288 available at: <https://www.bio-rad-antibodies.com/SDS/C12CA10288>

Regulatory

For research purposes only

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M350330:190307'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)