

Datasheet: C12CA BATCH NUMBER 156119

Description:	BABY RABBIT COMPLEMENT
Name:	BABY RABBIT COMPLEMENT
Format:	Serum
Product Type:	Serum
Quantity:	2 ml

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u> rad-antibodies.com/protocols.						
		Yes	No	Not Determined	Suggested Dilution		
	Functional Assays (1)	•					
	Immunoassay	•					
	Where this product has r	not been t	ested for ι	ise in a particular tech	nique this does not		
	necessarily exclude its u a guide only. It is recomm system using appropriate (1) This product is not It is preferable to dilute filtration in order to min	mended th e negative sold as s e the com	hat the use positive c terile but plement t	r titrates the product for controls. can be sterilized by f o a final working con	filtration if necessary.		
Product Form	Baby rabbit serum - lyop	hilised					
Reconstitution	Reconstitute with 2ml ice	e cold dist	illed water				
Preservative Stabilisers	None present						
Product Information	Baby rabbit complement complement for cytotoxic			n is intended for use a	s a source of rabbit		
Instructions For Use	Use within one hour of re	econstituti	on, keepin	g on ice throughout.			
References	1. Council of Europe. (19 of reagents and standard ISBN 92-871.224-4. 2. Edited by J.G.Ray. (19	disation of	complem	ent. European Health (Committee, Strasbourg,		

NIH-80-545.

3. Anderson, L.D. Jr *et al.* (1999) Enhancement of graft-versus-tumor activity and graft-versus-host disease by pretransplant immunization of allogeneic bone marrow donors with a recipient-derived tumor cell vaccine. <u>Cancer Res. 59 (7): 1525-30.</u>

4. Mason, J.C. *et al.* (2002) Statin-induced expression of decay-accelerating factor protects vascular endothelium against complement-mediated injury. <u>Circ Res. 91 (8)</u>: <u>696-703.</u>

5. Lidington, E.A. *et al.* (2000) Induction of decay-accelerating factor by thrombin through a protease-activated receptor 1 and protein kinase C-dependent pathway protects vascular endothelial cells from complement-mediated injury. <u>Blood. 96 (8): 2784-92.</u>

6. De clercq, L. *et al.* (1997) An anti-adipocyte monoclonal antibody is cytotoxic to porcine preadipocytes *in vitro* and depresses the development of pig adipose tissue. J Anim Sci. <u>75 (7): 1791-7.</u>

7. Hung, M.C. *et al.* (2011) The *Neisseria meningitidis* Macrophage Infectivity Potentiator Protein Induces Cross-Strain Serum Bactericidal Activity and Is a Potential Serogroup B Vaccine Candidate. <u>Infect Immun. 79: 3784-91.</u>

8. Lee, S.J. *et al.* (2012) Identification of a common immune signature in murine and human systemic Salmonellosis. <u>Proc Natl Acad Sci U S A. 109 (13): 4998-5003.</u>

9. Goh, Y.S. & MacLennan, C.A. (2013) Invasive African nontyphoidal Salmonella requires high levels of complement for cell-free antibody-dependent killing. <u>J Immunol Methods</u>. <u>387 (1-2): 121-9.</u>

10. Li, S.H. *et al.* (2004) C-reactive protein upregulates complement-inhibitory factors in endothelial cells. <u>Circulation. 109: 833-6.</u>

 Hyams, C. *et al.* (2010) *Streptococcus pneumoniae* resistance to complementmediated immunity is dependent on the capsular serotype. <u>Infect Immun. 78: 716-25.</u>
Newcombe, J. *et al.* (2004) Infection with an avirulent phoP mutant of *Neisseria meningitidis* confers broad cross-reactive immunity. Infect Immun. 72: 338-44.

13. Mason, J.C. *et al.* (2002) bFGF and VEGF synergistically enhance endothelial cytoprotection via decay-accelerating factor induction. <u>Am J Physiol Cell Physiol. 282:</u> <u>C578-87.</u>

14. Sancho, D. *et al.* (2006) CD69 targeting differentially affects the course of collageninduced arthritis. <u>J Leukoc Biol. 80: 1233-41.</u>

15. Hung MC *et al.* (2013) The adhesin complex protein (ACP) of *Neisseria meningitidis* is a new adhesin with vaccine potential. <u>MBio. 4 (2): pii: e00041-13.</u>

16. Goh YS *et al.* (2016) Bactericidal Immunity to *Salmonella* in Africans and Mechanisms Causing Its Failure in HIV Infection. <u>PLoS Negl Trop Dis. 10 (4): e0004604.</u>

17. Sawant S *et al.* (2016) Establishment of 3D Co-Culture Models from Different Stages of Human Tongue Tumorigenesis: Utility in Understanding Neoplastic Progression. <u>PLoS</u> <u>One. 11 (8): e0160615.</u>

18. Humbert MV *et al.* (2016) Vaccine Potential and Diversity of the Putative Cell Binding Factor (CBF, NMB0345/NEIS1825) Protein of *Neisseria meningitidis*. <u>PLoS One. 11 (8)</u>: <u>e0160403.</u>

19. Nganje, C.N. *et al.* (2019) PepN is a non-essential, cell wall-localized protein that contributes to neutrophil elastase-mediated killing of *Streptococcus pneumoniae*. <u>PLoS</u> <u>One. 14 (2): e0211632.</u>

20. Dierckx de Casterlé I *et al.* (2018) Reduction of myeloid-derived suppressor cells reinforces the anti-solid tumor effect of recipient leukocyte infusion in murine

			, .		ric Antigen Receptor T-Cell		
Storage		Prior to reconstitution store at +4°C. Following reconstitution store at +4°C for 1 hour or aliquot and store at -70°C for longer.					
		•	Should this product cor		eezing and thawing as this m tate we recommend		
Guarantee (Guaranteed until date of expiry. Please see product label.					
		Material Safety Datasheet documentation #10288 available at: https://www.bio-rad-antibodies.com/SDS/C12CA 10288					
Health A Informati	nd Safety ion	https://www.bio-rad-a			e at:		
	ion	https://www.bio-rad-a	ntibodies.com/SDS/C120		e at:		

Printed on 23 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint