

## Datasheet: C12CA

**BATCH NUMBER 171350**

<b>Description:</b>	BABY RABBIT COMPLEMENT
<b>Name:</b>	BABY RABBIT COMPLEMENT
<b>Format:</b>	Serum
<b>Product Type:</b>	Serum
<b>Quantity:</b>	2 ml

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Functional Assays (1)	▪			
Immunoassay	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

**(1) This product is not sold as sterile but can be sterilized by filtration if necessary. It is preferable to dilute the complement to a final working concentration before filtration in order to minimize loss of volume.**

**Product Form** Baby rabbit serum - lyophilized

**Reconstitution** Reconstitute with 2ml ice cold distilled water

**Preservative Stabilisers** None present

**Product Information** **Baby rabbit complement** serum preparation is intended for use as a source of rabbit complement for cytotoxicity assays.

**Instructions For Use** Use within one hour of reconstitution, keeping on ice throughout.

**References**

1. De clerq, L. *et al.* (1997) An anti-adipocyte monoclonal antibody is cytotoxic to porcine preadipocytes *in vitro* and depresses the development of pig adipose tissue. [J Anim Sci. 75 \(7\): 1791-7.](#)
2. Anderson, L.D. Jr *et al.* (1999) Enhancement of graft-versus-tumor activity and graft-

- versus-host disease by pretransplant immunization of allogeneic bone marrow donors with a recipient-derived tumor cell vaccine. [Cancer Res. 59 \(7\): 1525-30.](#)
3. Lidington, E.A. *et al.* (2000) Induction of decay-accelerating factor by thrombin through a protease-activated receptor 1 and protein kinase C-dependent pathway protects vascular endothelial cells from complement-mediated injury. [Blood. 96 \(8\): 2784-92.](#)
  4. Mason, J.C. *et al.* (2002) bFGF and VEGF synergistically enhance endothelial cytoprotection via decay-accelerating factor induction. [Am J Physiol Cell Physiol. 282: C578-87.](#)
  5. Mason, J.C. *et al.* (2002) Statin-induced expression of decay-accelerating factor protects vascular endothelium against complement-mediated injury. [Circ Res. 91 \(8\): 696-703.](#)
  6. Li, S.H. *et al.* (2004) C-reactive protein upregulates complement-inhibitory factors in endothelial cells. [Circulation. 109: 833-6.](#)
  7. Newcombe, J. *et al.* (2004) Infection with an avirulent phoP mutant of *Neisseria meningitidis* confers broad cross-reactive immunity. [Infect Immun. 72: 338-44.](#)
  8. Sancho, D. *et al.* (2006) CD69 targeting differentially affects the course of collagen-induced arthritis. [J Leukoc Biol. 80: 1233-41.](#)
  9. Hyams, C. *et al.* (2010) *Streptococcus pneumoniae* resistance to complement-mediated immunity is dependent on the capsular serotype. [Infect Immun. 78: 716-25.](#)
  10. Hung, M.C. *et al.* (2011) The *Neisseria meningitidis* Macrophage Infectivity Potentiator Protein Induces Cross-Strain Serum Bactericidal Activity and Is a Potential Serogroup B Vaccine Candidate. [Infect Immun. 79: 3784-91.](#)
  11. Lee, S.J. *et al.* (2012) Identification of a common immune signature in murine and human systemic Salmonellosis. [Proc Natl Acad Sci U S A. 109 \(13\): 4998-5003.](#)
  12. Hung MC *et al.* (2013) The adhesin complex protein (ACP) of *Neisseria meningitidis* is a new adhesin with vaccine potential. [MBio. 4 \(2\): pii: e00041-13.](#)
  13. Goh, Y.S. & MacLennan, C.A. (2013) Invasive African nontyphoidal Salmonella requires high levels of complement for cell-free antibody-dependent killing. [J Immunol Methods. 387 \(1-2\): 121-9.](#)
  14. Goh YS *et al.* (2016) Bactericidal Immunity to *Salmonella* in Africans and Mechanisms Causing Its Failure in HIV Infection. [PLoS Negl Trop Dis. 10 \(4\): e0004604.](#)
  15. Humbert MV *et al.* (2016) Vaccine Potential and Diversity of the Putative Cell Binding Factor (CBF, NMB0345/NEIS1825) Protein of *Neisseria meningitidis*. [PLoS One. 11 \(8\): e0160403.](#)
  16. Dierckx de Casterlé I *et al.* (2018) Reduction of myeloid-derived suppressor cells reinforces the anti-solid tumor effect of recipient leukocyte infusion in murine neuroblastoma-bearing allogeneic bone marrow chimeras. [Cancer Immunol Immunother. 67 \(4\): 589-603.](#)
  17. Valton, J. *et al.* (2018) A Versatile Safeguard for Chimeric Antigen Receptor T-Cell Immunotherapies. [Sci Rep. 8 \(1\): 8972.](#)
  18. Dierckx de Casterlé, I. *et al.* (2018) Reduction of myeloid-derived suppressor cells reinforces the anti-solid tumor effect of recipient leukocyte infusion in murine neuroblastoma-bearing allogeneic bone marrow chimeras. [Cancer Immunol Immunother. 67 \(4\): 589-603.](#)
  19. Nganje, C.N. *et al.* (2019) PepN is a non-essential, cell wall-localized protein that contributes to neutrophil elastase-mediated killing of *Streptococcus pneumoniae*. [PLoS One. 14 \(2\): e0211632.](#)

20. Cuesta-Mateos, C. *et al.* (2020) CCR7 as a novel therapeutic target in t-cell PROLYMPHOCYTIC leukemia [Biomarker Research.8, 54.](#)
21. Mosti, L. *et al.* (2021) Targeted multi-epitope switching enables straightforward positive/negative selection of CAR T cells. [Gene Ther. 28 \(9\): 602-12.](#)
22. Olivera-Ardid, S. *et al.* (2023) Removal of natural anti-αGal antibodies elicits protective immunity against Gram-negative bacterial infections. [Front Immunol. 14: 1232924.](#)
23. Flaugnatti, N. *et al.* (2025) Capsular polysaccharide restrains type VI secretion in *Acinetobacter baumannii*. [Elife. 14Jan 03 \[Epub ahead of print\].](#)

---

**Storage**

This product is shipped at ambient temperature.  
Prior to reconstitution store at +4°C. Following reconstitution store at +4°C for 1 hour or aliquot and store at -70°C for longer.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the product. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee**

Guaranteed until date of expiry. Please see product label.

---

**Health And Safety Information**

Material Safety Datasheet documentation #10288 available at: <https://www.bio-rad-antibodies.com/SDS/C12CA>

---

**Regulatory**

For research purposes only

---

**Product inquiries:** [www.bio-rad-antibodies.com/technical-support](http://www.bio-rad-antibodies.com/technical-support)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](http://bio-rad-antibodies.com/datasheets)

'M439339:250523'

**Printed on 08 Mar 2026**