

Datasheet: DC002

Description:	MOUSE ANTI HUMAN CD3:FITC/CD4:RPE
Specificity:	CD3/CD4
Format:	FITC/RPE
Product Type:	Dual Color Reagent
Clone:	UCHT1/RPA-T4
Isotype:	Cocktail
Quantity:	50 TESTS

Product Details

RRID AB_1125251

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Antibody Isotypes FITC reagent: IgG1 (MOUSE)
RPE reagent: IgG1 (MOUSE)

Target Species Human

Product Form Dual Colour combination consisting of FITC conjugated and RPE conjugated monoclonal antibodies mixed in optimal ratio - lyophilised.

Reconstitution Reconstitute with 0.5 ml distilled water

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
	RPE 488nm laser	496	578
	RPE 561nm laser	546	578

Buffer Solution Phosphate buffered saline

Preservative 0.09% Sodium Azide
Stabilisers 1% Bovine Serum Albumin
5% Sucrose

**External Database
Links**

UniProt:

[P04234](#) [Related reagents](#)
[P07766](#) [Related reagents](#)
[P01730](#) [Related reagents](#)
[P09693](#) [Related reagents](#)

Entrez Gene:

[915](#) CD3D [Related reagents](#)
[916](#) CD3E [Related reagents](#)
[917](#) CD3G [Related reagents](#)
[920](#) CD4 [Related reagents](#)

Synonyms

T3D, T3E, T3G

Specificity

UCHT1 recognises the human CD3 antigen, which is expressed by all mature T cells.

RPA-T4 recognises the human CD4 antigen, a 56 kDa glycoprotein expressed by the helper-inducer subset of T cells. CD4 is also weakly expressed by monocytes.

DC002 may be used to simultaneously identify CD3 and CD4 positive cells, and to identify populations of cells that co-express these antigens.

Flow Cytometry

Use 10ul to label 100ul whole blood.

References

1. Beverley, P.C. & Callard, R.E. (1981) Distinctive functional characteristics of human "T" lymphocytes defined by E rosetting or a monoclonal anti-T cell antibody. [Eur J Immunol. 11 \(4\): 329-34.](#)
2. Parnes, J.R. (1989) Molecular biology and function of CD4 and CD8. [Adv Immunol. 44: 265-311.](#)

Storage

Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of reconstitution.

**Health And Safety
Information**

Material Safety Datasheet documentation #10075 available at:
10075: <https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf>

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1:FITC/MOUSE IgG1:RPE NEGATIVE CONTROL \(DC012\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M335407:181204'

Printed on 20 May 2019

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)