

Datasheet: DC047

BATCH NUMBER 154471

Description:	ANTI DOG CD3:FITC/CD8:RPE
Specificity:	CD3/CD8
Format:	FITC/RPE
Product Type:	Dual Color Reagent
Clone:	CA17.2A12 / YCATE55.9
Isotype:	Cocktail
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own systems using appropriate negative/positive controls.

Antibody Isotypes	FITC reagent: IgG1 (MOUSE) RPE reagent: IgG1 (RAT)		
Target Species	Dog		
Product Form	Dual Colour combination consisting of FITC conjugated and RPE conjugated monoclonal antibodies mixed in optimal ratio - lyophilised.		
Reconstitution	Reconstitute with 1.0 ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
	RPE 488nm laser	496	578
	RPE 561nm laser	546	578
Buffer Solution	Phosphate buffered saline		

Preservative	0.09% Sodium Azide
Stabilisers	1% Bovine Serum Albumin
	5% Sucrose

External Database Links

UniProt:

[P27597](#) [Related reagents](#)
[P33706](#) [Related reagents](#)

Entrez Gene:

[442981](#) CD3E [Related reagents](#)
[403157](#) CD8A [Related reagents](#)

RRID AB_324122

Specificity **Anti Canine CD3 / CD8 dual color reagent** is a dual colour reagent recognising canine CD3 and CD8 cell surface antigens. Clone CA17.2A12 recognises canine CD3, a T cell differentiation antigen expressed on mature T lymphocytes.

Clone YCATE 55.9 recognises canine CD8, which is expressed by cytotoxic T cells. YCATE 55.9 binds to CD8 alpha.

Flow Cytometry Use 10ul of the suggested working dilution to label 10^6 cells in 100ul.

References 1. Sakai, M *et al.* (2006) Phenotypic analysis of hepatic T lymphocytes in a dog with chronic hepatitis. [J Vet Med Sci. 68:1219-1221](#)

Storage Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #20487 available at: <https://www.bio-rad-antibodies.com/SDS/DC04720487>

Regulatory For research purposes only

North & South America Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M375221:210104'

