

Datasheet: HCA053A

Description:	HUMAN ANTI Ki67
Specificity:	Ki67
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	AbD02531
Isotype:	HuCAL/mouse IgG1 Fc
Quantity:	50 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen	▪			
Immunohistology - Paraffin (1)	▪			1/100 - 1/500
ELISA	▪			2 µg/ml
Western Blotting	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

Target Species	Human
Species Cross Reactivity	Based on sequence similarity, is expected to react with: Bovine, Macaque, Dog, Chimpanzee, Rhesus Monkey N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Chimeric human-mouse IgG1 antibody selected from the HuCAL® phage display library and expressed in a human cell line. The antibody has variable regions of human origin and an Fc portion (including CH1 and CL domains) from mouse IgG1. It can be detected by anti mouse Fc specific secondary antibodies. This antibody is supplied lyophilized.
Reconstitution	Reconstitute with 0.5 ml distilled water
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin

Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	Peptide derived from the human Ki67 protein, sequence GFKELFQTPG, coupled via a C-terminal cysteine to carrier proteins
External Database Links	<p>UniProt: P46013 Related reagents</p> <p>Entrez Gene: 4288 MKI67 Related reagents</p>
Specificity	<p>Human anti Ki67 antibody, clone AbD02531 recognizes the Ki67 cell-cycle associated protein. Ki67 is expressed in proliferating cells but not in quiescent cells. Expression of this antigen occurs preferentially during late G1, S, G2, and M phases of the cell cycle, while in cells in G0 phase the antigen cannot be detected. Consequently, Ki-67 antigen expression is used in tumor pathology to detect proliferating cells in neoplastic diseases. In cultured cells, Ki-67 is expressed in the nucleolus of interphase cells. The Ki67 gene contains 15 exons. The Ki67 repeat region, within which there is a 22-amino acid Ki67 motif, is encoded by exon 13. The shorter isoform lacks exon 7. Northern blot analysis reveals multiple transcripts ranging from approximately 8.9 to 12.5 kb in proliferating but not quiescent cells. Immunoblot analysis shows expression of 320 and 359 kDa proteins. Antisense oligonucleotides inhibit cellular proliferation in a dose-dependent manner, suggesting that Ki67 protein expression may be an absolute requirement for cell proliferation.</p> <p>Within cells Ki67 is predominantly localized in the G1 phase in the perinucleolar region, in the later phases it is also detected throughout the nuclear interior, being predominantly localized in the nuclear matrix. In mitosis, it is present on all chromosomes.</p>
ELISA	Due to the presence of bovine serum albumin (BSA), this antibody is unsuitable for use as a capture reagent in sandwich ELISA applications. This product is also available without BSA, please enquire .
Histology Positive Control Tissue	Human tonsil
References	<ol style="list-style-type: none"> Jarutat, T. <i>et al.</i> (2006) Isolation and comparative characterization of Ki-67 equivalent antibodies from the HuCAL phage display library. Biol. Chem. 387: 995-1003. Liu, S.K. <i>et al.</i> (2011) Delta-like ligand 4-notch blockade and tumor radiation response. J Natl Cancer Inst. 103 (23): 1778-98.
Storage	<p>Prior to reconstitution store at +4°C.</p> <p>After reconstitution store at +4°C or at -20°C if preferred.</p> <p>Storage in frost-free freezers is not recommended.</p>
Shelf Life	12 months from date of reconstitution
Acknowledgements	Sold under license of U.S. Patents 6753136, 7785859 and 8273688 and corresponding patents. This antibody was developed by Bio-Rad, Zeppelinstr. 4, 82178 Puchheim, Germany.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf

Licensed Use For *in vitro* research purposes only, unless otherwise specified in writing by Bio-Rad.

Regulatory For research purposes only

Technical Advice Recommended protocols and further information about HuCAL recombinant antibody technology can be found in the [HuCAL Antibodies Technical Manual](#)

Related Products

Recommended Secondary Antibodies

Goat Anti Human IgG F(ab')₂ (0500-0099...) [HRP](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M319646:180726'

Printed on 23 Oct 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)