

## Datasheet: HCA117

<b>Description:</b>	HUMAN ANTI Ki67
<b>Specificity:</b>	Ki67
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	AbD02815
<b>Isotype:</b>	HuCAL Fab bivalent
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin (1)	▪			
ELISA	▪			
Immunoprecipitation			▪	
Western Blotting			▪	
Functional Assays			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

**(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.**

<b>Target Species</b>	Human
<b>Product Form</b>	A bivalent human recombinant Fab selected from the HuCAL® phage display library, expressed in <i>E. coli</i> . This Fab fragment is bivalent by dimerization of the bacterial alkaline phosphatase fusion protein. The antibody is tagged with a DYKDDDDK tag and a HIS-tag (HHHHHH) at the C-terminus of the antibody heavy chain. This antibody is supplied liquid.
<b>Preparation</b>	metal chelate affinity chromatography
<b>Buffer Solution</b>	Phosphate buffered saline

<b>Preservative Stabilisers</b>	0.01% Thiomersal
<b>Approx. Protein Concentrations</b>	Total protein concentration 0.5 mg/ml
<b>Immunogen</b>	Peptide derived from the human Ki67 protein, sequence GFKELFQTPG, coupled via a C-terminal cysteine to carrier proteins
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P46013</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">4288</a>    MKI67    <a href="#">Related reagents</a></p>
<b>RRID</b>	AB_2142235
<b>Specificity</b>	<p><b>Human anti Ki67 antibody, clone AbD02815</b> recognizes the sequence GFKELFQ, which can be found eight times on the human Ki67 protein. Ki67 antigen is expressed in proliferating cells but not in quiescent cells. Expression of this antigen occurs preferentially during late G1, S, G2, and M phases of the cell cycle, while in cells in G0 phase the antigen cannot be detected. Consequently, Ki67 antigen expression is used in tumor pathology to detect proliferating cells in neoplastic diseases. In cultured cells, Ki67 is expressed in the nucleolus of interphase cells.</p> <p>The Ki67 gene contains 15 exons. The Ki67 repeat region, within which there is a 22-amino acid Ki67 motif, is encoded by exon 13. The shorter isoform lacks exon 7. Northern blot analysis revealed multiple transcripts ranging from approximately 8.9 to 12.5 kb in proliferating but not quiescent cells. Immunoblot analysis showed expression of 320 and 359 kDa proteins. Sequence analysis predicted that the short-lived 2,896- and 3,256-amino acid protein isoforms contain potential nuclear targeting signals, over 200 potential phosphorylation sites, 19 N-myristoylation sites, 3 amidation sites, and numerous PEST sites. Antisense oligonucleotides inhibited cellular proliferation in a dose-dependent manner, suggesting that Ki67 protein expression may be an absolute requirement for cell proliferation.</p>
<b>Histology Positive Control Tissue</b>	Human tonsil
<b>Storage</b>	<p>Store at +4°C or at -20°C if preferred.  Storage in frost-free freezers is not recommended.  This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
<b>Guarantee</b>	12 months from date of despatch
<b>Acknowledgements</b>	Sold under license of U.S. Patents 6753136, 7785859 and 8273688 and corresponding

patents. This antibody was developed by Bio-Rad, Zeppelinstr. 4, 82178 Puchheim, Germany.

His-tag is a registered trademark of EMD Biosciences.

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10094 available at: 10094: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10094.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10094.pdf</a>
<b>Licensed Use</b>	For in vitro research purposes only, unless otherwise specified in writing by Bio-Rad.
<b>Regulatory</b>	For research purposes only
<b>Technical Advice</b>	Recommended protocols and further information about HuCAL recombinant antibody technology can be found in the <a href="#">HuCAL Antibodies Technical Manual</a>

---

## Related Products

### Recommended Secondary Antibodies

Mouse Anti Synthetic Peptide HISTIDINE TAG (MCA5995...) [HRP](#)

Rat Anti Synthetic Peptide DYKDDDDK TAG (MCA4764...) [HRP](#)

**North & South America** Tel: +1 800 265 7376  
Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

From March 15, 2021, we will no longer supply printed datasheets with our products.  
Look out for updates on how to access your digital version at [bio-rad-antibodies.com](http://bio-rad-antibodies.com)

'M360873:200124'

**Printed on 09 Feb 2021**

---

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)