

Datasheet: HCA150A

Description:	HUMAN ANTI HUMAN PINK1
Specificity:	PINK1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	AbD11408
lsotype:	HuCAL Fab monovalent
Quantity:	50 µg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further						
	information. For general	information. For general protocol recommendations, please visit <u>www.bio-</u>					
	rad-antibodies.com/protocols.						
		Yes	No	Not Determined	Suggested Dilution		
	Flow Cytometry						
	Immunohistology - Frozen	•			1/50		
	Immunohistology - Paraffin (1)	•			1/50		
	ELISA						
	Immunoprecipitation						
	Western Blotting	•			2 ug/ml		
	Functional Assays			•			
	a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls. (1)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections.Sodium citrate buffer pH 6.0 is recommended for this purpose.						
Target Species	Human						
Product Form	A monovalent human recombinant Fab (lambda light chain) selected from the HuCAL® phage display library, expressed in <i>E. coli</i> The antibody is tagged with a V5 tag and a double extended Strep-tag at the C-terminus of the antibody heavy chain. This antibody is supplied lyophilized.						
Reconstitution	Reconstitute with 0.5 ml	distilled v	vater				
Preparation	StrepTactin affinity chrom	atograph	ıy				

Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide (NaN ₃)
Stabilisers	1% Bovine Serum Albumin
Approx. Protein	
Concentrations	Ig concentration 0.1 mg/ml after reconstitution
Concontrationo	
Immunogen	Peptide corresponding to amino acids 321 – 511 of human PINK1
External Database	
Links	UniProt:
Links	Q9BXM7 Related reagents
	QBDAMT Related reagents
	Entrez Gene:
	65018 PINK1 Related reagents
RRID	AB_10845154
	-
Specificity	Human anti Human PINK1 antibody, clone AbD11408 recognizes human
	PTEN-induced putative kinase protein 1 (PINK1), a 581 aa single pass membrane protein
	belonging the serine/threonine protein kinase.
	PINK1 protects against mitochondrial dysfunction during cellular stress, possibly by
	phosphorylating mitochondrial proteins. PINK1 is highly expressed in tissues such as the
	heart, skeletal muscle and testis. Lower expression levels are detected in the brain,
	·
	placenta, liver, kidney, pancreas, prostate, ovary and small intestine.
	placenta, liver, kidney, pancreas, prostate, ovary and small intestine.
	placenta, liver, kidney, pancreas, prostate, ovary and small intestine. Defects in PINK1 are the cause of autosomal recessive early-onset Parkinson's disease 6
	placenta, liver, kidney, pancreas, prostate, ovary and small intestine.
	placenta, liver, kidney, pancreas, prostate, ovary and small intestine. Defects in PINK1 are the cause of autosomal recessive early-onset Parkinson's disease 6
	placenta, liver, kidney, pancreas, prostate, ovary and small intestine. Defects in PINK1 are the cause of autosomal recessive early-onset Parkinson's disease 6 (<u>PARK6</u>), a condition characterized by the early onset of Parkinsonian symptoms including
Histology Positive	placenta, liver, kidney, pancreas, prostate, ovary and small intestine. Defects in PINK1 are the cause of autosomal recessive early-onset Parkinson's disease 6 (<u>PARK6</u>), a condition characterized by the early onset of Parkinsonian symptoms including tremor, bradykinesia, possible hyperreflexia and dementia (Valente <i>et al.</i> 2004).
Histology Positive Control Tissue	placenta, liver, kidney, pancreas, prostate, ovary and small intestine. Defects in PINK1 are the cause of autosomal recessive early-onset Parkinson's disease 6 (<u>PARK6</u>), a condition characterized by the early onset of Parkinsonian symptoms including
	placenta, liver, kidney, pancreas, prostate, ovary and small intestine. Defects in PINK1 are the cause of autosomal recessive early-onset Parkinson's disease 6 (<u>PARK6</u>), a condition characterized by the early onset of Parkinsonian symptoms including tremor, bradykinesia, possible hyperreflexia and dementia (Valente <i>et al.</i> 2004).
Control Tissue	placenta, liver, kidney, pancreas, prostate, ovary and small intestine. Defects in PINK1 are the cause of autosomal recessive early-onset Parkinson's disease 6 (PARK6), a condition characterized by the early onset of Parkinsonian symptoms including tremor, bradykinesia, possible hyperreflexia and dementia (Valente <i>et al.</i> 2004). Human testis
	placenta, liver, kidney, pancreas, prostate, ovary and small intestine. Defects in PINK1 are the cause of autosomal recessive early-onset Parkinson's disease 6 (<u>PARK6</u>), a condition characterized by the early onset of Parkinsonian symptoms including tremor, bradykinesia, possible hyperreflexia and dementia (Valente <i>et al.</i> 2004).
Control Tissue Western Blotting	 placenta, liver, kidney, pancreas, prostate, ovary and small intestine. Defects in PINK1 are the cause of autosomal recessive early-onset Parkinson's disease 6 (PARK6), a condition characterized by the early onset of Parkinsonian symptoms including tremor, bradykinesia, possible hyperreflexia and dementia (Valente <i>et al.</i> 2004). Human testis HCA150A detects a band of approximately 70kDa in human brain cell lysates
Control Tissue	 placenta, liver, kidney, pancreas, prostate, ovary and small intestine. Defects in PINK1 are the cause of autosomal recessive early-onset Parkinson's disease 6 (PARK6), a condition characterized by the early onset of Parkinsonian symptoms including tremor, bradykinesia, possible hyperreflexia and dementia (Valente <i>et al.</i> 2004). Human testis HCA150A detects a band of approximately 70kDa in human brain cell lysates 1. Marzetti, E. <i>et al.</i> (2016) Altered mitochondrial quality control signaling in muscle of old
Control Tissue Western Blotting	 placenta, liver, kidney, pancreas, prostate, ovary and small intestine. Defects in PINK1 are the cause of autosomal recessive early-onset Parkinson's disease 6 (PARK6), a condition characterized by the early onset of Parkinsonian symptoms including tremor, bradykinesia, possible hyperreflexia and dementia (Valente <i>et al.</i> 2004). Human testis HCA150A detects a band of approximately 70kDa in human brain cell lysates 1. Marzetti, E. <i>et al.</i> (2016) Altered mitochondrial quality control signaling in muscle of old gastric cancer patients with cachexia. Exp Gerontol. Nov 12. pii: S0531-5565(16)30259-5.
Control Tissue Western Blotting	 placenta, liver, kidney, pancreas, prostate, ovary and small intestine. Defects in PINK1 are the cause of autosomal recessive early-onset Parkinson's disease 6 (PARK6), a condition characterized by the early onset of Parkinsonian symptoms including tremor, bradykinesia, possible hyperreflexia and dementia (Valente <i>et al.</i> 2004). Human testis HCA150A detects a band of approximately 70kDa in human brain cell lysates 1. Marzetti, E. <i>et al.</i> (2016) Altered mitochondrial quality control signaling in muscle of old
Control Tissue Western Blotting	 placenta, liver, kidney, pancreas, prostate, ovary and small intestine. Defects in PINK1 are the cause of autosomal recessive early-onset Parkinson's disease 6 (PARK6), a condition characterized by the early onset of Parkinsonian symptoms including tremor, bradykinesia, possible hyperreflexia and dementia (Valente <i>et al.</i> 2004). Human testis HCA150A detects a band of approximately 70kDa in human brain cell lysates 1. Marzetti, E. <i>et al.</i> (2016) Altered mitochondrial quality control signaling in muscle of old gastric cancer patients with cachexia. Exp Gerontol. Nov 12. pii: S0531-5565(16)30259-5.
Control Tissue Western Blotting	 placenta, liver, kidney, pancreas, prostate, ovary and small intestine. Defects in PINK1 are the cause of autosomal recessive early-onset Parkinson's disease 6 (PARK6), a condition characterized by the early onset of Parkinsonian symptoms including tremor, bradykinesia, possible hyperreflexia and dementia (Valente <i>et al.</i> 2004). Human testis HCA150A detects a band of approximately 70kDa in human brain cell lysates 1. Marzetti, E. <i>et al.</i> (2016) Altered mitochondrial quality control signaling in muscle of old gastric cancer patients with cachexia. Exp Gerontol. Nov 12. pii: S0531-5565(16)30259-5.
Control Tissue Western Blotting References	placenta, liver, kidney, pancreas, prostate, ovary and small intestine. Defects in PINK1 are the cause of autosomal recessive early-onset Parkinson's disease 6 (PARK6), a condition characterized by the early onset of Parkinsonian symptoms including tremor, bradykinesia, possible hyperreflexia and dementia (Valente et al. 2004). Human testis HCA150A detects a band of approximately 70kDa in human brain cell lysates 1. Marzetti, E. et al. (2016) Altered mitochondrial quality control signaling in muscle of old gastric cancer patients with cachexia. Exp Gerontol. Nov 12. pii: S0531-5565(16)30259-5. [Epub ahead of print] Prior to reconstitution store at +4°C.
Control Tissue Western Blotting References	 placenta, liver, kidney, pancreas, prostate, ovary and small intestine. Defects in PINK1 are the cause of autosomal recessive early-onset Parkinson's disease 6 (PARK6), a condition characterized by the early onset of Parkinsonian symptoms including tremor, bradykinesia, possible hyperreflexia and dementia (Valente et al. 2004). Human testis HCA150A detects a band of approximately 70kDa in human brain cell lysates 1. Marzetti, E. et al. (2016) Altered mitochondrial quality control signaling in muscle of old gastric cancer patients with cachexia. Exp Gerontol. Nov 12. pii: S0531-5565(16)30259-5. [Epub ahead of print] Prior to reconstitution store at +4°C. After reconstitution store at +4°C or at -20°C if preferred.
Control Tissue Western Blotting References	 placenta, liver, kidney, pancreas, prostate, ovary and small intestine. Defects in PINK1 are the cause of autosomal recessive early-onset Parkinson's disease 6 (PARK6), a condition characterized by the early onset of Parkinsonian symptoms including tremor, bradykinesia, possible hyperreflexia and dementia (Valente <i>et al.</i> 2004). Human testis HCA150A detects a band of approximately 70kDa in human brain cell lysates 1. Marzetti, E. <i>et al.</i> (2016) Altered mitochondrial quality control signaling in muscle of old gastric cancer patients with cachexia. Exp Gerontol. Nov 12. pii: S0531-5565(16)30259-5. [Epub ahead of print] Prior to reconstitution store at +4°C. After reconstitution store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored
Control Tissue Western Blotting References	 placenta, liver, kidney, pancreas, prostate, ovary and small intestine. Defects in PINK1 are the cause of autosomal recessive early-onset Parkinson's disease 6 (PARK6), a condition characterized by the early onset of Parkinsonian symptoms including tremor, bradykinesia, possible hyperreflexia and dementia (Valente <i>et al.</i> 2004). Human testis HCA150A detects a band of approximately 70kDa in human brain cell lysates 1. Marzetti, E. <i>et al.</i> (2016) Altered mitochondrial quality control signaling in muscle of old gastric cancer patients with cachexia. Exp Gerontol. Nov 12. pii: S0531-5565(16)30259-5. [Epub ahead of print] Prior to reconstitution store at +4°C. After reconstitution store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should
Control Tissue Western Blotting References	 placenta, liver, kidney, pancreas, prostate, ovary and small intestine. Defects in PINK1 are the cause of autosomal recessive early-onset Parkinson's disease 6 (PARK6), a condition characterized by the early onset of Parkinsonian symptoms including tremor, bradykinesia, possible hyperreflexia and dementia (Valente <i>et al.</i> 2004). Human testis HCA150A detects a band of approximately 70kDa in human brain cell lysates 1. Marzetti, E. <i>et al.</i> (2016) Altered mitochondrial quality control signaling in muscle of old gastric cancer patients with cachexia. Exp Gerontol. Nov 12. pii: S0531-5565(16)30259-5. [Epub ahead of print] Prior to reconstitution store at +4°C. After reconstitution store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored
Control Tissue Western Blotting References	 placenta, liver, kidney, pancreas, prostate, ovary and small intestine. Defects in PINK1 are the cause of autosomal recessive early-onset Parkinson's disease 6 (PARK6), a condition characterized by the early onset of Parkinsonian symptoms including tremor, bradykinesia, possible hyperreflexia and dementia (Valente <i>et al.</i> 2004). Human testis HCA150A detects a band of approximately 70kDa in human brain cell lysates 1. Marzetti, E. <i>et al.</i> (2016) Altered mitochondrial quality control signaling in muscle of old gastric cancer patients with cachexia. Exp Gerontol. Nov 12, pii: S0531-5565(16)30259-5. [Epub ahead of print] Prior to reconstitution store at +4°C. After reconstitution store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should
Control Tissue Western Blotting References	 placenta, liver, kidney, pancreas, prostate, ovary and small intestine. Defects in PINK1 are the cause of autosomal recessive early-onset Parkinson's disease 6 (PARK6), a condition characterized by the early onset of Parkinsonian symptoms including tremor, bradykinesia, possible hyperreflexia and dementia (Valente <i>et al.</i> 2004). Human testis HCA150A detects a band of approximately 70kDa in human brain cell lysates 1. Marzetti, E. <i>et al.</i> (2016) Altered mitochondrial quality control signaling in muscle of old gastric cancer patients with cachexia. Exp Gerontol. Nov 12. pii: S0531-5565(16)30259-5. [Epub ahead of print] Prior to reconstitution store at +4°C. After reconstitution store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should

Acknowledgements	Sold under license of U.S. Patents 6753136, 7785859 and 8273688 and corresponding patents. This antibody was developed by Bio-Rad, Zeppelinstr. 4, 82178 Puchheim, Germany.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf</u>
Regulatory	For research purposes only
Technical Advice	Recommended protocols and further information about HuCAL recombinant antibody technology can be found in the <u>HuCAL Antibodies Technical Manual</u>

Related Products

Recommended Secondary Antibodies

Mouse Anti Synthetic Peptide STREP-TAG CLASSIC (MCA2489...) HRP

Mouse Anti Viral V5-TAG (MCA1360...)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@	bio-rad.com	Email: antibody_sales_uk@bio	o-rad.com	Email: antibody_sales_de@bio-rad.com
			no longer supply printed of to access your digital vers 'M333971:181127'		the second se
			Printed on 10 Feb 2021		

Alk. Phos., Purified

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint