

## Datasheet: HCA355

Description:	ANTI MONO-ADP-RIBOSE
Specificity:	MONO-ADP-RIBOSE
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	AbD33205
lsotype:	lgG
Quantity:	0.1 mg

## **Product Details**

Applications	This product has been re derived from testing with communications from the information. For general	tions or personal dicated for further					
	rad-antibodies.com/proto						
		Yes	No	Not Determined	Suggested Dilution		
	Western Blotting	•			2 ug/ml		
	Immunofluorescence				1/100		
	Where this product has not been tested for use in a particular technique this does not						
	necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.						
Target Species	Protein/peptide tag						
Product Form	Human/Rabbit IgG chim phage display library an liquid.	•					
Preparation	Purified recombinant Ig0 mammalian cell line.	G prepared	l by affinit	y chromatography on I	Protein A from a		
Source	НКВ-11						
Buffer Solution	Phosphate buffered salir	ne					
Preservative Stabilisers	0.09% Sodium Azide (N	aN <sub>3</sub> )					
Carrier Free	Yes						

Approx. Protein Concentrations	Total protein concentration 0.5 mg/ml
Immunogen	ARTKQTARKS(ADPr)TGGKAC
Specificity	Anti mono-ADP-ribose antibody, clone AbD33205, recognizes mono-adenosine diphosphate (ADP)-ribosylation. ADP-ribosylation is a reversible post-translational modification that occurs in multicellular organisms as well as some lower unicellular eukaryotes, but is absent in prokaryotes and yeast (Bürkle 2005). ADP ribosylation has been shown to play critical roles in many physiological and pathological processes, including bacterial pathogenesis and signaling and metabolism to control chromatin-related processes including transcription and DNA repair (Bonfiglio <i>et al.</i> 2020, Bütepage <i>et al.</i> 2015).
	Members of the ADP-ribosyltransferase (ART) superfamily of proteins including the poly(ADP-ribose) polymerases (PARPs) subfamily, catalyze the transfer of ADP-ribose from nicotinamide adenine dinucleotide (NAD+) onto substrate protein via N-, O-, or S-glycosidic linkages. These transferases are able to transfer a single ADP-ribose residue to their substrate proteins, in a process known as mono-ADP-ribosylation ( <u>Bütepage <i>et al.</i></u> 2015).
	This clone AbD33205 is a mono-selective ADP-ribose antibody which was generated using an H3S10ADPr peptide as the antigen. This clone is specific for mono-ADPr and recognizes Ser-mono-ADPr. Another mono-ADPr antibody ( <u>HCA354</u> , AbD33204) also specifically recognizes mono-ADP-ribose, however it has a preference for mono-ADP-riboses other than Ser-mono-ADPr, with a preference for ubiquitin Arg-ADPr and mono-ADPr catalyzed by other PARPs.
	Anti mono-ADP-ribose antibody, clone AbD33205, has been used in immunofluorescence staining of H2O2-treated U2OS cells to showed strong nuclear staining, relating to HPF1 and PARP1, but not PARP2. This staining was abolished upon treatment with the PARP inhibitors (Bonfiglio <i>et al.</i> 2020).
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Acknowledgements	This product and/or its use is covered by claims of U.S. patents, and/or pending U.S. and non-U.S. patent applications owned by or under license to Bio-Rad Laboratories, Inc. See bio-rad.com/en-us/trademarks for details.

Information <u>https://www.bio-rad-ar</u> 10040		antibodies.com/SDS/HCA	<u>.355</u>		
Regulator	ry	For research purpose	es only		
North & South America	Tel: +1 800 265 7376 <b>Worldwide</b> Fax: +1 919 878 3751		Tel: +44 (0)1865 852 700 <b>Europe</b> Fax: +44 (0)1865 852 739		Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50
	Email: antibody_s	sales_us@bio-rad.com	Email: antibody_sales_uk@bio	o-rad.com	Email: antibody_sales_de@bio-rad.com

## Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint