

## Datasheet: HCA355

<b>Description:</b>	ANTI MONO-ADP-RIBOSE
<b>Specificity:</b>	MONO-ADP-RIBOSE
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	AbD33205
<b>Isotype:</b>	IgG
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			2 ug/ml
Immunofluorescence	▪			1/100

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Protein/peptide tag
<b>Product Form</b>	Human/Rabbit IgG chimera (rabbit CH2 and CH3) antibody selected from the HuCAL phage display library and expressed in a human cell line. This antibody is supplied as a liquid.
<b>Preparation</b>	Purified recombinant IgG prepared by affinity chromatography on Protein A from a mammalian cell line.
<b>Source</b>	HKB-11
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.01% Thiomersal
<b>Carrier Free</b>	Yes

<b>Approx. Protein Concentrations</b>	Total protein concentration 0.5 mg/ml
<b>Immunogen</b>	ARTKQTARKS(ADPr)TGGKAC
<b>Specificity</b>	<p><b>Anti mono-ADP-ribose antibody, clone AbD33205</b>, recognizes mono-adenosine diphosphate (ADP)-ribosylation. ADP-ribosylation is a reversible post-translational modification that occurs in multicellular organisms as well as some lower unicellular eukaryotes, but is absent in prokaryotes and yeast (<a href="#">Bürkle 2005</a>). ADP ribosylation has been shown to play critical roles in many physiological and pathological processes, including bacterial pathogenesis and signaling and metabolism to control chromatin-related processes including transcription and DNA repair (<a href="#">Bonfiglio et al. 2020</a>, <a href="#">Bütepage et al. 2015</a>).</p> <p>Members of the ADP-ribosyltransferase (ART) superfamily of proteins including the poly(ADP-ribose) polymerases (PARPs) subfamily, catalyze the transfer of ADP-ribose from nicotinamide adenine dinucleotide (NAD<sup>+</sup>) onto substrate protein via N-, O-, or S-glycosidic linkages. These transferases are able to transfer a single ADP-ribose residue to their substrate proteins, in a process known as mono-ADP-ribosylation (<a href="#">Bütepage et al. 2015</a>).</p> <p>This clone AbD33205 is a mono-selective ADP-ribose antibody which was generated using an H3S10ADPr peptide as the antigen. This clone is specific for mono-ADPr and recognizes Ser-mono-ADPr. Another mono-ADPr antibody (<a href="#">HCA354</a>, AbD33204) also specifically recognizes mono-ADP-ribose, however it has a preference for mono-ADP-riboses other than Ser-mono-ADPr, with a preference for ubiquitin Arg-ADPr and mono-ADPr catalyzed by other PARPs.</p> <p>Anti mono-ADP-ribose antibody, clone AbD33205, has been used in immunofluorescence staining of H<sub>2</sub>O<sub>2</sub>-treated U2OS cells to showed strong nuclear staining, relating to HPF1 and PARP1, but not PARP2. This staining was abolished upon treatment with the PARP inhibitors (<a href="#">Bonfiglio et al. 2020</a>).</p>
<b>Storage</b>	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
<b>Guarantee</b>	12 months from date of despatch
<b>Acknowledgements</b>	This product and/or its use is covered by claims of U.S. patents, and/or pending U.S. and non-U.S. patent applications owned by or under license to Bio-Rad Laboratories, Inc. See <a href="http://bio-rad.com/en-us/trademarks">bio-rad.com/en-us/trademarks</a> for details.

---

**Health And Safety Information**      Material Safety Datasheet documentation #10094 available at:  
10094: <https://www.bio-rad-antibodies.com/uploads/MSDS/10094.pdf>

---

**Regulatory**                      For research purposes only

---

**North & South America**      Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M416897:230302'

**Printed on 02 Mar 2023**

---

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)