

Datasheet: ICT913 BATCH NUMBER 159083

Description:	FAM FLICA™ CASPASE-9 KIT
Name:	CASPASE-9
Format:	FAM (Green Fluorescence)
Product Type:	Kits
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			
Immunofluorescence				

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FAM	494	520
Product Information	The FAM FLICA Casp	•	ck and easy method t

Test Principle

Caspase FLICA kits measure apoptosis by detecting active caspases in whole, living cells. These kits do not work by using antibodies or as an ELISA. Instead, their methodology is based on a unique cell-permeable and non-cytotoxic reagent called the Fluorochrome Inhibitor of Caspases (FLICA). The FLICA reagent contains a caspase inhibitor sequence linked to a green (Carboxyfluorescein, FAM) fluorescent probe.

The Caspase FLICA Kits are suitable for cells in suspension and adherent cells from many species including mammalian, insect and yeast. Different cell types, e.g. HeLa, primary neurons, macrophages and lymphocytes have also been successfully studied with these kits.

This kit can be used with a flow cytometer, fluorescence microscope or a fluorescence

plate reader using black microtitre plates.

Reagents In The Kit

4 vials of FAM-LEHD-FMK FLICA Reagent - lyophilized

10x Apoptosis Wash Buffer, 60 mL

Fixative, 6 mL

Propidium Iodide, 1 mL Hoechst 33342, 1 mL

Instructions For Use

Instructions for use can be found at www.bio-rad-antibodies.com/uploads/IFU/ICT913.pdf

References

1. Brugnon, F. et al. (2010) Apoptosis and meiotic segregation in ejaculated sperm from Robertsonian translocation carrier patients. <u>Hum Reprod. 25 (7): 1631-42.</u>

2. Rébé, C. et al. (2007) Caspase-8 prevents sustained activation of NF-kappaB in monocytes undergoing macrophagic differentiation. <u>Blood. 109 (4): 1442-50.</u>

3. Vacher, P. et al. (2015) Localized Store-Operated Calcium Influx Represses CD95-Dependent Apoptotic Effects of Rituximab in Non-Hodgkin B Lymphomas. J

Immunol. 195 (5): 2207-15.

Storage

Store the unopened kit and each unopened component at +4°C until the expiration date. Once reconstituted with DMSO, use FLICA reagent immediately, or store at -20°C for 6 months protected from light and thawed no more than twice during that time.

Guarantee

Guaranteed until date of expiry. Please see product label.

Acknowledgements

FLICA™ is a trademark of Immunochemistry Technologies, LLC.

Health And Safety Information

Material Safety Datasheet documentation #20285 #10471 #10498 #10476 #10477

available at:

https://www.bio-rad-antibodies.com/SDS/ICT913

FAM-LEHD-FMK FLICA Reagent (20285) 10x Apoptosis Wash Buffer (10471)

Fixative (10498)

Hoechst 33342 (10476) Propidium Iodide (10477)

Regulatory

For research purposes only

North & South Tel: +1 800 265 7376

America

Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21

Email: antibody_sales_us@bio-rad.com

Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M374934:201208'

Printed on 29 Feb 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint