

## Datasheet: ICT9152

**BATCH NUMBER 166518**

<b>Description:</b>	GREEN CATHEPSIN B KIT
<b>Name:</b>	CATHEPSIN B
<b>Format:</b>	Rhodamine 110-(RR)2
<b>Product Type:</b>	Kits
<b>Quantity:</b>	100 TESTS

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Refer to Instruction for Use
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	Rhodamine110-(RR)2	525	535

#### Product Information

**Green Cathepsin B Kit** enables the quantitation and monitoring of intracellular cathepsin activity over time in vitro. The Rhodamine 110 Cathepsin B substrate reagent is a non-cytotoxic and membrane permeant substrate that fluoresces green upon cleavage by active cathepsin enzymes.

#### Test Principle

Rhodamine 110 Cathepsin B substrate utilizes the photostable green fluorophore, rhodamine 110. Rhodamine 110 cathepsin B substrate is comprised of rhodamine 110 coupled to two copies of the amino acid sequence, arginine-arginine (RR), which is the preferential target sequence for cathepsin B. When bi-substituted via amide linkage to two cathepsin B target peptide sequences, rhodamine110 is nonfluorescent. Following enzymatic cleavage at one or both arginine (R) amide linkage sites, the mono and non-substituted rhodamine 110 fluorophores generate green fluorescence when excited at 500 nm.

To use the Green Cathepsin B Assay, simply add the Rhodamine 110 Cathepsin B

substrate [R110-(RR)<sub>2</sub>] directly to the cell culture media (or 1X Cellular Assay Buffer), incubate, and analyze. Because R110-(RR)<sub>2</sub> is cell-permeant, it easily penetrates the cell membrane and the membranes of the internal cellular organelles - no lysis or permeabilization steps are required. R110-(RR)<sub>2</sub> will enter the cell in a non-fluorescent state. If cathepsin enzymes are active, they will cleave off the two arginine-arginine cathepsin B targeting sequences and allow the rhodamine 110 fluorophore to become fluorescent upon excitation. By varying the duration and concentration of exposure to the R110-(RR)<sub>2</sub> substrate, a picture can be obtained of the relative abundance and intracellular location of cathepsin enzymatic activity. Positive cells will fluoresce green, while negative cells will exhibit very low levels of background green fluorescence. There is no interference from pro-cathepsins forms of the enzymes. If the treatment or experimental condition stimulates cathepsin activity, cells containing elevated levels of cathepsin activity will appear brighter green than cells with lower levels of cathepsin activity.

---

**Reagents In The Kit** 4 vial of Rhodamine110-(RR)<sub>2</sub> substrate - lyophilized  
10X Cellular Assay Buffer, 60 ml  
Hoechst Stain, 1 ml

---

**Instructions For Use** Instructions for use can be found at <https://www.bio-rad-antibodies.com/static/uploads/ifu/ict9151-2.pdf>

---

**Storage** MULTIPLE STORAGE CONDITIONS APPLY ON ARRIVAL. Store the unopened kit (and each unopened component) according to the storage instructions on each component label. Store the Rhodamine110-(RR)<sub>2</sub> substrate at -20°C. Once reconstituted in DMSO, use the Rhodamine110-(RR)<sub>2</sub> substrate immediately or aliquot and store at -20°C for 6 months, protected from light. Avoid repeated freezing and thawing

---

**Guarantee** Guaranteed until date of expiry. Please see product label.

---

**Health And Safety Information** Material Safety Datasheet documentation #20428 #20429 #10476 available at: <https://www.bio-rad-antibodies.com/SDS/ICT9152>  
Rhodamine 110-(RR)<sub>2</sub> Substrate (20428)  
10X Cellular Assay Buffer (20429)  
Hoechst Stain (10476)

---

**Regulatory** For research purposes only

---

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M360122:191030'

Printed on 18 Jan 2024