

Datasheet: ICT917 BATCH NUMBER 157164

Description:	SR FLICA™ POLY CASPASE KIT
Name:	POLY CASPASE
Format:	Sulforhodamine (Red Fluorescence)
Product Type:	Kits
Quantity:	100 TESTS

Product Details

	derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u>					
	rad-antibodies.com/protocols. Yes No Not Determined Suggested Dilution					
	Flow Cytometry Immunofluorescence	•				
	a guide only. It is reco	mmended that the us	er titrates the product f	for use in their own		
Mars For/For	system using appropri	iate negative/positive	controls.			
Max Ex/Em	•	iate negative/positive				
Max Ex/Em Product Information	system using appropri Fluorophore Sulforhodamine The SR FLICA Poly C caspases in apoptotic caspase 1, caspase 3	Excitation Max (nm) 565 Caspase Kit uses a c cells. This FLICA rea , caspase 4, caspase	controls. Emission Max (nm)	to analyze active g activated caspases: e 7, caspase 8 and		

cells. These kits do not work by using antibodies or as an ELISA. Instead, their methodology is based on a unique cell-permeable and non-cytotoxic reagent called the Fluorochrome Inhibitor of Caspases (FLICA). The FLICA™ reagent contains a caspase inhibitor sequence linked to a red (Sulforhodamine, SR) fluorescent probe.

The Caspase FLICA[™] Kits are suitable for cells in suspension and adherent cells from many species including mammalian, insect and yeast. Different cell types, e.g. HeLa, primary neurons, macrophages and lymphocytes have also been successfully studied with these kits.

	This kit can be used with a fluorescence microscope or a fluorescence plate reader using black microtitre plates.
Reagents In The Kit	4 vials of SR-VAD-FMK FLICA™ Reagent - lyophilized 10X Wash Buffer, 60 mL Fixative, 6 mL Hoechst Stain, 1 mL
Instructions For Use	Instructions for use can be found at www.bio-rad-antibodies.com/uploads/IFU/ICT917.pdf
References	 Soni, C. <i>et al.</i> (2015) Distinct and synergistic roles of FcγRIIB deficiency and 129 strain- derived SLAM family proteins in the development of spontaneous germinal centers and autoimmunity. <u>J Autoimmun. 63: 31-46.</u> Wong, E.B. <i>et al.</i> (2015) B cell-intrinsic CD84 and Ly108 maintain germinal center B cell tolerance. <u>J Immunol. 194 (9): 4130-43.</u>
Storage	Store the unopened kit (and each unopened component) at +4°C until the expiration date. Protect the FLICA™ reagent from light at all times.
	Once reconstituted, the 150X FLICA™ stock should be stored at -20°C protected from light.
Guarantee	Guaranteed until date of expiry. Please see product label.
Acknowledgements	FLICA™ is a trademark of Immunochemistry Technologies, LLC.
Health And Safety Information	Material Safety Datasheet documentation #20289 #10471 #10498 #10476 available at: <u>https://www.bio-rad-antibodies.com/SDS/ICT917</u> SR-VAD-FMK FLICA™ Reagent (20289) 10X Wash Buffer (10471) Fixative (10498) Hoechst Stain (10476)
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

FAM FLICA™ CASPASE-3/7 KIT (ICT094)FAM FLICA™ CASPASE-1 KIT (ICT098)FAM FLICA™ CASPASE-6 KIT (ICT096)FAM FLICA™ CASPASE-8 KIT (ICT910)FAM FLICA™ CASPASE-9 KIT (ICT913)FAM FLICA™ CASPASE-10 KIT (ICT923)

North & South	Tel: +1 800 265 7376	Worldwide
America	Fax: +1 919 878 3751	
	Email: antibody_sales_us@bio-rad.	.com

Tel: +44 (0)1865 852 700 **Europe** Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

Printed on 29 Feb 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint