

Datasheet: ICT944

MITOCHONDRIAL PERMEABILITY TRANSITION KIT
MITOCHONDRIAL PERMEABILITY TRANSITION:MitoPT™ JC-1 KIT
JC-1 Dye (Dual Green/Red Fluorescence)
Kits
400 TESTS

## **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="https://www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			
Immunofluorescence	-			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

## **Product Information**

**The MitoPT JC-1 kit** uses a quick and easy staining method to clearly differentiate between non-apoptotic and apoptotic cells through mitochondrial functionality.

### **Test Principle**

An early indication of apoptosis involves a collapse in the electrochemical gradient across the mitochondrial membrane. Loss of mitochondrial membrane potential can be detected by a unique fluorescent cationic dye known as JC-1 (5,5',6,6'-tetrachloro-1,1',3,3'-tetraethylbenzamidazolocarbocyanin iodide) that has been incorporated into the MitoPT JC-1 kit.

The MitoPT JC-1 reagent easily penetrates cells and enters the mitochondria. It aggregates in the mitochondria of non-apoptotic cells and fluoresces red, whilst in apoptotic cells it diffuses throughout the cell. Once dispersed, the reagent assumes a monomeric form and fluoresces green. This allows an easy distinction between non-apoptotic red fluorescent cells and apoptotic green fluorescent cells which can be read with a flow cytometer, fluorescence microscope or a fluorescence plate reader using black microtitre plates.

## Reagents In The Kit

MitoPT JC-1 Reagent, 400 tests Two bottles 10x Assay Buffer, 125 mL CCCP (50 mM in DMSO), 600 ul.

Instructions For Use	Instructions for use can be found at <a href="https://www.bio-rad-antibodies.com/uploads/IFU/ICT944.pg">www.bio-rad-antibodies.com/uploads/IFU/ICT944.pg</a>
References	1. Jasek, E. et al. (2012) Effect of histone deacetylase inhibitors trichostatin A and valpr
	acid on etoposide-induced apoptosis in leukemia cells. <u>Anticancer Res. 32 (7): 2791-9.</u>
	2. Wei, X. et al. (2014) In Vitro Comparative Effect of Three Novel Borate Bioglasses or
	the Behaviors of Osteoblastic MC3T3-E1 Cells <u>J Mat Sci Technol. 30 (10): 979-83.</u>
	3. Gizak, A. et al. (2019) Targeting a moonlighting function of aldolase induces apoptos
	in cancer cells. Cell Death Dis. 10 (10): 712.
	4. Lasota, M. et al. (2022) Property of a new Prussian blue-bounded iron complex able
	peroxidate non-saturated fatty acids with a tendency to create conditions that may
	encourage ferroptosis. <u>J Physiol Pharmacol. 73 (5): 673-86.</u>
	5. Gizak, A. <i>et al.</i> (2021) A novel remitting leukodystrophy associated with a variant in
	FBP2. Brain Commun. 3 (2): fcab036.
	FBP2. <u>Brain Commun. 3 (2). Icabo36.</u>
Storage	
Storage	Store the unopened kit and each unopened component at -20°C until the expiration dat
Storage	
Storage	Once opened, some components may be stored at 2-8°C until the expiration date. CCC
Storage	Once opened, some components may be stored at 2-8°C until the expiration date. CCC should be stored frozen. Once reconstituted with DMSO, dilute and use MitoPT reagent
Storage	Once opened, some components may be stored at 2-8°C until the expiration date. CCC should be stored frozen. Once reconstituted with DMSO, dilute and use MitoPT reagen immediately, or store at ≤-20°C for 12 months protected from light and thawed no more
Storage	Store the unopened kit and each unopened component at -20°C until the expiration date. Once opened, some components may be stored at 2-8°C until the expiration date. CCC should be stored frozen. Once reconstituted with DMSO, dilute and use MitoPT reagent immediately, or store at ≤-20°C for 12 months protected from light and thawed no more than twice.
	Once opened, some components may be stored at 2-8°C until the expiration date. CCC should be stored frozen. Once reconstituted with DMSO, dilute and use MitoPT reagent immediately, or store at ≤-20°C for 12 months protected from light and thawed no more
Storage  Guarantee  Acknowledgements	Once opened, some components may be stored at 2-8°C until the expiration date. CCC should be stored frozen. Once reconstituted with DMSO, dilute and use MitoPT reagent immediately, or store at ≤-20°C for 12 months protected from light and thawed no more than twice.
Guarantee Acknowledgements	Once opened, some components may be stored at 2-8°C until the expiration date. CCC should be stored frozen. Once reconstituted with DMSO, dilute and use MitoPT reagent immediately, or store at ≤-20°C for 12 months protected from light and thawed no more than twice.  Guaranteed until date of expiry. Please see product label.  MitoPT is a trademark of Immunochemistry Technologies, LLC.
Guarantee Acknowledgements Health And Safety	Once opened, some components may be stored at 2-8°C until the expiration date. CCC should be stored frozen. Once reconstituted with DMSO, dilute and use MitoPT reagent immediately, or store at ≤-20°C for 12 months protected from light and thawed no more than twice.  Guaranteed until date of expiry. Please see product label.  MitoPT is a trademark of Immunochemistry Technologies, LLC.  Material Safety Datasheet documentation #20291 #10474 #10479 available at:
Guarantee	Once opened, some components may be stored at 2-8°C until the expiration date. CCC should be stored frozen. Once reconstituted with DMSO, dilute and use MitoPT reagent immediately, or store at ≤-20°C for 12 months protected from light and thawed no more than twice.  Guaranteed until date of expiry. Please see product label.  MitoPT is a trademark of Immunochemistry Technologies, LLC.  Material Safety Datasheet documentation #20291 #10474 #10479 available at: https://www.bio-rad-antibodies.com/SDS/ICT944
Guarantee Acknowledgements Health And Safety	Once opened, some components may be stored at 2-8°C until the expiration date. CCC should be stored frozen. Once reconstituted with DMSO, dilute and use MitoPT reagent immediately, or store at ≤-20°C for 12 months protected from light and thawed no more than twice.  Guaranteed until date of expiry. Please see product label.  MitoPT is a trademark of Immunochemistry Technologies, LLC.  Material Safety Datasheet documentation #20291 #10474 #10479 available at: https://www.bio-rad-antibodies.com/SDS/ICT944  MitoPT JC-1 Reagent (20291)
Guarantee Acknowledgements Health And Safety	Once opened, some components may be stored at 2-8°C until the expiration date. CCC should be stored frozen. Once reconstituted with DMSO, dilute and use MitoPT reagent immediately, or store at ≤-20°C for 12 months protected from light and thawed no more than twice.  Guaranteed until date of expiry. Please see product label.  MitoPT is a trademark of Immunochemistry Technologies, LLC.  Material Safety Datasheet documentation #20291 #10474 #10479 available at: https://www.bio-rad-antibodies.com/SDS/ICT944  MitoPT JC-1 Reagent (20291) 10x Assay Buffer (10474)
Guarantee Acknowledgements Health And Safety	Once opened, some components may be stored at 2-8°C until the expiration date. CCC should be stored frozen. Once reconstituted with DMSO, dilute and use MitoPT reagent immediately, or store at ≤-20°C for 12 months protected from light and thawed no more than twice.  Guaranteed until date of expiry. Please see product label.  MitoPT is a trademark of Immunochemistry Technologies, LLC.  Material Safety Datasheet documentation #20291 #10474 #10479 available at: https://www.bio-rad-antibodies.com/SDS/ICT944  MitoPT JC-1 Reagent (20291)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M404843:220906'

# Printed on 25 Jun 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint