

## Datasheet: LNK002P

Description:	LYNX RAPID H	RP ANTIBO	DY CONJ	UGATION KIT		
Name:	HRP CONJUGATION KIT Kit Conjugation Kit 3 CONJUGATIONS for 400µg antibody					
Format:						
Product Type:						
Quantity:						
roduct Details						
Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .					
	Conjugation	Yes	Νο	Not Determined	Suggested Dilution	
Product Information	LYNX Rapid HRP Antibody Conjugation Kit® enables the rapid conjugation of a pre-prepared lyophilized mixture containing Horseradish peroxidase (HRP) label to an antibody or protein. Activation of proprietary reagents within the antibody-label solution results in directional covalent bonding of HRP to the antibody. The LYNX Rapid Conjugation kit® can be used to label small quantities of antibody/protein at near neutral pH, allowing a high conjugation efficiency with 100%					
Reagents In The Kit	antibody recovery. 3 Vials of 100ug LYNX 1 Vial LYNX Modifier r	( lyophilized	_		,	
	1 Vial LYNX Quenche	0				
Preparing The Antibody	The following buffer solutions are recommended for preparing the antibody:					
	10-50mM amine-free buffer (e.g HEPES, MES, MOPS and phosphate) pH range 6.5-8.5, although moderate concentrations of Tris buffer (<20mM) may be tolerated.					
	although moderate co	ncentrations			, · · ·	

Sodium azide is an irreversible inhibitor of HRP and therefore should be avoided.

The amount of antibody used for labeling ideally should correspond to molar ratios between 1:4 and 1:1 Ab to HRP. Taking account of the molecular weights (160,000 versus 40,000), this means for that for 100 $\mu$ g HRP you need to add between 100-400 $\mu$ g of antibody. For optimal results the antibody volume should be up to 100 $\mu$ l, at a concentration range of 0.5-5.0mg/ml.

Instructions For Use 1. To the antibody sample add 1ul of the Modifier reagent for every 10ul of antibody and mix gently.

2. Pipette the mixed antibody-modifier sample directly onto the LYNX lyophilized mix and gently pipette up and down twice to resuspend.

3. Replace cap onto vial and incubate at room temperature (20-25°C) for 3 hours, or overnight if preferred.

4. After incubation, add 1ul of Quencher reagent for every 10ul of antibody used. Leave to stand for 30 minutes before use.

1. Bondzio, A. et al. (2011) Identification of differentially expressed proteins in ruminal
epithelium in response to a concentrate-supplemented diet. Am J Physiol Gastrointest
Liver Physiol. 301 (2): G260-8.
2. Lichtmannegger, J. <i>et al.</i> (2016) Methanobactin reverses acute liver failure in a rat
model of Wilson disease. <u>J Clin Invest. 126 (7): 2721-35.</u>
3. Sasson, S.C. et al. (2021) Identification of neutralising pembrolizumab anti-drug
antibodies in patients with melanoma. <u>Sci Rep. 11 (1): 19253.</u>
4. Rosadas, C. et al. (2022) Detection and quantification of antibody to SARS CoV 2
receptor binding domain provides enhanced sensitivity, specificity and utility. <u>J Virol</u>
<u>Methods. 302: 114475.</u>
5. Khan, M. et al. (2022) Simple, sensitive, specific self-sampling assay secures
SARS-CoV-2 antibody signals in sero-prevalence and post-vaccine studies. Sci Rep. 12
<u>(1): 1885.</u>
6. Saraban, K. et al. (2023) Hybrid immunity from SARS-CoV-2 infection and mRNA
BNT162b2 vaccine among Thai school-aged children. Vaccine X. 15: 100414.
7. Peraile, I. <i>et al.</i> (2023) STUDY OF THE REUSABILITY AND STABILITY OF NYLON
NANOFIBRES AS AN ANTIBODY IMMOBILISATION SURFACE. Beilstein Arch. 9 Oct
[Epub ahead of print].
This kit contains lyophilized hygroscopic components that are moisture-sensitive. This kit
is shipped under ambient conditions with silica packets to avoid exposure to moisture. On
receipt, Bio-Rad recommend that the kit is stored at -20°C and protected from moisture.
Storage in frost-free freezers is not recommended. This product should be stored
undiluted. Avoid repeated freezing and thawing. Before opening, allow the components to
reach room temperature to minimize condensation.

Health And		12 months from date o	of despatch		
Information	'n	•	46)		#10548 available at:
Licensed L		Kingdom patent number purchase of this product right (without the right use the product to make purchaser cannot sell of data made using this p purchasing licenses for at. Endeavour House,	er 2446088 and associa ct conveys to the buyer to resell repackage or fi ke conjugates for resear or otherwise transfer thi product, or its componer r diagnostic and other u Langford Business Parl	ated internation the limited, in urther sublice rch and deve s product, or nts to a third uses may be k, Langford L	atent protected under United onal patent applications. The non exclusive non-transferable ense) under these patents to elopment purposes only. The r its components, or materials of party. Further information on obtained by contacting Bio-Rad ane, Kidlington, Oxon. OX5 antibodies@bio-rad.com
Regulatory	y	For research purposes	only		
<b>merica</b> F	Tel: +1 800 265 73 Fax: +1 919 878 37 Email: antibody_sa		Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio	Europe -rad.com	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
	tch/lot specific	datasheet for this produ	ct, please use our online s	search tool at:	
To find a bat			'M390795:211004'		bio-rad-antibodies.com/datasheets