

## Datasheet: LNK033APC BATCH NUMBER 164890

Description:	LYNX RAPID APC ANTIBODY CONJUGATION KIT		
Name:	APC CONJUGATION KIT		
Format:	Kit		
Product Type:	Conjugation Kit		
Quantity:	1 CONJUGATION for 1mg antibody		

## **Product Details**

Applications	This product has been red derived from testing with communications from the information. For general rad-antibodies.com/proto Conjugation We recommend that for e ratio.	in our labo e originator protocol re <u>ocols</u> . Yes	ratories, p rs. Please commenc <b>No</b>	peer-reviewed publicat refer to references in dations, please visit <u>w</u> <b>Not Determined</b>	tions or personal dicated for further <u>ww.bio-</u> Suggested Dilution	
Product Information	LYNX Rapid APC Antibody Conjugation Kit® enables the rapid conjugation of a pre-prepared lyophilized mixture containing Allophycocyanin (APC) label to an antibody or protein. Activation of proprietary reagents within the antibody-label solution results in directional covalent bonding of APC to the antibody. The LYNX Rapid Conjugation kit® can be used to label small quantities of antibody/protein at near neutral pH, allowing a high conjugation efficiency with 100% antibody recovery.					
Reagents In The Kit	1 Vial of 1mg LYNX lyop 1 Vial LYNX Modifier rea 1 Vial LYNX Quencher re	gent	C mix			
Preparing The Antibody	The following buffer solutions are recommended for preparing the antibody:					
	10-50mM amine-free buffer (e.g HEPES, MES, MOPS and phosphate) pH range 6.5-8.5, although moderate concentrations of Tris buffer (<20mM) may be tolerated.					
	If possible, avoid buffers containing nucleophilic components such as prima amines and thiols (e.g. thiomersal/thimerosal) since they may react with LYN					

chemicals. Azide (0.02-0.1%), EDTA, up to 50% Glycerol and common non-buffering salts and sugars have little or no effect on conjugation efficiency. For optimal results the antibody should be at a concentration of 1mg/ml, with a maximum volume of 1ml and a maximum antibody amount of 1mg. Antibody at a concentration of greater than 1mg/ml requires dilution. Antibody below 1mg/ml can still be used as long as the maximum volume is not exceeded. Using less than the recommended amount of antibody may result in unbound label, but this will be removed during subsequent application wash steps. Antibody below 0.5mg/ml should be concentrated before use with the kit. Instructions For Use 1. To the antibody sample add 1ul of the Modifier reagent for every 10ul of antibody and mix gently. 2. Pipette the mixed antibody-modifier sample directly onto the LYNX lyophilized mix and gently pipette up and down twice to resuspend. 3. Replace cap onto vial and incubate in the dark at room temperature (20-25°C) for 3 hours, or overnight if preferred. 4. After incubation, add 1ul of Quencher reagent for every 10ul of antibody used. Leave to stand for 30 minutes before use. References 1. Wang, Y. et al. (2010) Local host response to chlamydial urethral infection in male guinea pigs. Infect Immun.78: 1670-81. 2. Lacy, H.M. et al. (2011) Essential Role for Neutrophils in Pathogenesis and Adaptive Immunity in Chlamydia caviae Ocular Infections. Infect Immun. 79: 1889-97 3. Paget, C. et al. (2012) Interleukin-22 is produced by invariant natural killer T lymphocytes during influenza A virus infection: potential role in protection against lung epithelial damage. J Biol Chem. 287: 8816-29. 4. Seliger, C. et al. (2011) A rapid high-precision flow cytometry based technique for total white blood cell counting in chickens. Vet Immunol Immunopathol. 145: 86-99. 5. Fu, Y. et al. (2014) Development of a FACS-based assay for evaluating antiviral potency of compound in dengue infected peripheral blood mononuclear cells. J Virol Methods. 196: 18-24. 6. TraxImayr, M.W. et al. (2014) Construction of pH-sensitive Her2-binding IgG1-Fc by directed evolution. Biotechnol J. 9: 1013-22. 7. Wielgosz, M.M. et al. (2015) Generation of a lentiviral vector producer cell clone for human Wiskott-Aldrich syndrome gene therapy. Mol Ther Methods Clin Dev. 2: 14063. 8. Hofer, C.C. et al. (2015) Infection of mice with influenza A/WSN/33 (H1N1) virus alters alveolar type II cell phenotype. Am J Physiol Lung Cell Mol Physiol. 308 (7): L628-38. 9. Poh, C.M. et al. (2014) Damage to the blood-brain barrier during experimental cerebral malaria results from synergistic effects of CD8+ T cells with different specificities. Infect Immun. 82: 4854-64. 10. Hasenhindl, C. et al. (2014) Creating stable stem regions for loop elongation in Fcabs - insights from combining yeast surface display, in silico loop reconstruction and molecular

Regulatory	For research purposes only						
	at. Endeavour House, Langford Business Park, Langford Lane, Kidlington, Oxon. OX5 1GE UNITED KINGDOM. Tel: +44 1865 852 700. E-mail: <u>antibodies@bio-rad.com</u>						
	purchasing licenses for diagnostic and other uses may be obtained by contacting Bio-Ra						
	data made using this product, or its components to a third party. Further information on						
	use the product to make conjugates for research and development purposes only. The purchaser cannot sell or otherwise transfer this product, or its components, or materials						
	right (without the right to resell repackage or further sublicense) under these patents to						
	purchase of this product conveys to the buyer the limited, non exclusive non-transferable						
	Kingdom patent number 2446088 and associated international patent applications. The						
Licensed Use	These products and the methodology of conjugation are patent protected under United						
	Quencher Reagent (10548)						
	Lyophilized APC Mix (10532) Modifier Reagent (10546)						
Information	https://www.bio-rad-antibodies.com/SDS/LNK033APC						
Health And Saf							
Guarantee	12 months from date of despatch						
	reach room temperature to minimize condensation.						
	undiluted. Avoid repeated freezing and thawing. Before opening, allow the components t						
	Storage in frost-free freezers is not recommended. This product should be stored						
	is shipped under ambient conditions with silica packets to avoid exposure to moisture. O receipt, Bio-Rad recommend that the kit is stored at -20°C and protected from moisture.						
Storage	This kit contains lyophilized hygroscopic components that are moisture-sensitive. This ki						
	and humoral immunity in pigs. <u>Virol J. 20 (1): 181.</u>						
	16. Haach, V. <i>et al.</i> (2023) A polyvalent virosomal influenza vaccine induces broad cellul						
	<ol> <li>Jax, E. <i>et al.</i> (2023) Evaluating Effects of AIV Infection Status on Ducks Using a Flow Cytometry-Based Differential Blood Count. <u>Microbiol Spectr. 11 (4): e0435122.</u></li> </ol>						
	<u>10406387221077969.</u>						
	the equine systemic inflammatory response syndrome. <u>J Vet Diagn Invest.</u>						
	14. Theuerkauf, K. <i>et al.</i> (2022) Activated platelets and platelet-leukocyte aggregates in						
	<u>32-43.</u>						
	13. Hercher, D. <i>et al.</i> (2020) Motor and sensory Schwann cell phenotype commitment is diminished by extracorporeal shockwave treatment <i>in vitro</i> <u>J Peripher Nerv Syst. 25 (1)</u>						
	improves schwann cell adhesion and proliferation. <u>J Mater Sci Mater Med. 27 (12): 188.</u>						
	12. Schuh, C.M. et al. (2016) Covalent binding of placental derived proteins to silk fibroir						
	cancer. <u>BMC Cancer. 16 (1): 154.</u>						
	11. Ward, S.T. <i>et al.</i> (2016) Evaluation of serum and tissue levels of VAP-1 in colorectal						
	dynamics simulations. <u>Biochim Biophys Acta.1844: 1530-40.</u>						

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

Email: antibody\_sales\_uk@bio-rad.com

Email: antibody\_sales\_de@bio-rad.com

Email: antibody\_sales\_us@bio-rad.com

'M390399:210910'

## Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint