

Datasheet: LNK061F BATCH NUMBER 168707

Description:	LYNX RAPID FLUORESCEIN ANTIBODY CONJUGATION KIT						
Name:	FLUORESCEIN CONJUGATION KIT						
Format:	Kit						
Product Type:	Conjugation Kit						
Quantity:	3 CONJUGATIONS for 200µg antibody						
Product Details							
Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .						
	Yes No Not Determined Suggested Dilution						
	Conjugation						
Product Information	LYNX Rapid Fluorescein Antibody Conjugation Kit ® enables the rapid conjugation of pre-prepared lyophilized mixture containing Fluorescein label to an antibody or protein. Activation of proprietary reagents within the antibody-label solution results in directional covalent bonding of Fluorescein to the antibody.						
	The LYNX Rapid Conjugation kit® can be used to label small quantities of antibody/protein at near neutral pH, allowing a high conjugation efficiency with 100% antibody recovery.						
Reagents In The Kit	3 Vials LYNX lyophilized Fluorescein mix 1 Vial LYNX Modifier reagent 1 Vial LYNX Quencher reagent						
Preparing The Antibody	The following buffer solutions are recommended for preparing the antibody:						
	10-50mM amine-free buffer (e.g HEPES, MES, MOPS and phosphate) pH range 6.5-8.5, although moderate concentrations of Tris buffer (<20mM) may be tolerated.						
	If possible, avoid buffers containing nucleophilic components such as primary amines and thiols (e.g. thiomersal/thimerosal) since they may react with LYNX						

chemicals. Azide (0.02-0.1%), EDTA and common non-buffering salts and sugars have little or no effect on conjugation efficiency.

It is recommended that 100-200ug antibody be used in each labelling reaction. For
optimal results the antibody should be at a concentration of 1mg/ml, with a maximum
volume of 100ul and a recommended antibody amount of 100ug. A maximum of 200ug of
antibody can be used to obtain good quality conjugates as long as the maximum
conjugation volume of 100ul is not exceeded. Antibody below 1mg/ml can still be used as
long as the maximum volume is not exceeded. Using less than the recommended amount
of antibody may result in unbound label, but this will be removed during subsequent
application wash steps. Antibody below 0.5mg/ml should be concentrated before use with
the kit.

Instructions For Use	1. To the antibody sample add 1ul of the Modifier reagent for every 10ul of antibody and mix gently.				
	2. Pipette the mixed antibody-modifier sample directly onto the LYNX lyophilized mix and gently pipette up and down twice to resuspend.				
	3. Replace cap onto vial and incubate in the dark at room temperature (20-25 ^o C) for 3 hours or overnight, if preferred.				
	4. After incubation, add 1ul of Quencher reagent for every 10ul of antibody used. Leave to stand for 30 minutes before use.				
References	1. Walker, E. <i>et al.</i> (2014) Microscopic Detection of Quenched Activity-Based Optical Imaging Probes Using an Antibody Detection System: Localizing Protease Activity. <u>Mol</u> <u>Imaging Biol. 16: 608-18.</u>				
	2. Mahata, B. <i>et al.</i> (2014) Single-cell RNA sequencing reveals T helper cells synthesizing steroids de novo to contribute to immune homeostasis. <u>Cell Rep. 7: 1130-42.</u>				
	3. Gawronska-Kozak, B. <i>et al.</i> (2021) Dermal White Adipose Tissue (dWAT) Is Regulated				
	by Foxn1 and Hif-1 α during the Early Phase of Skin Wound Healing. Int J Mol Sci. 23 (1): 257.				
	 4. Jax, E. <i>et al.</i> (2023) Evaluating Effects of AIV Infection Status on Ducks Using a Flow Cytometry-Based Differential Blood Count. <u>Microbiol Spectr. 11 (4): e0435122.</u> 				
Storage	This kit contains lyophilized hygroscopic components that are moisture-sensitive. This kit is shipped under ambient conditions with silica packets to avoid exposure to moisture. On receipt, Bio-Rad recommend that the kit is stored at -20°C and protected from moisture. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing. Before opening, allow the components to reach room temperature to minimize condensation.				
Guarantee	12 months from date of despatch				
Health And Safety Information	Material Safety Datasheet documentation #10534 #10546 #10549 available at: https://www.bio-rad-antibodies.com/SDS/LNK061F				

		Modifier I	ed Fluoresce Reagent (105 er Reagent (1			
Licensed	l Use	Kingdom purchase right (with use the p purchase data mac purchasin at. Endea	patent numb of this produ- hout the right product to ma er cannot sell de using this p ng licenses for avour House,	her 2446088 and associated conveys to the buyer to resell repackage or for the conjugates for resear or otherwise transfer the product, or its compone or diagnostic and other the Langford Business Par	ated internation the limited, rurther sublic rch and deve is product, o nts to a third uses may be k, Langford I	patent protected under United ional patent applications. The non exclusive non-transferable cense) under these patents to elopment purposes only. The rr its components, or materials or l party. Further information on obtained by contacting Bio-Rad, Lane, Kidlington, Oxon. OX5 antibodies@bio-rad.com
Regulatory		For resea	arch purpose:			
North & South America	Tel: +1 800 265 7 Fax: +1 919 878 : Email: antibody_s	3751	Worldwide rad.com	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bic	Europe o-rad.com	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
To find a b	atch/lot specifi	c datashee	et for this produ	uct, please use our online 'M390401:210910'	search tool at	: bio-rad-antibodies.com/datasheets
				Printed on 26 Jun 2024		

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint