

## Datasheet: LNK062F BATCH NUMBER 169153

Description:	LYNX RAPID	FLUORESCE	IN ANTIB	ODY CONJUGATION	I KIT		
Name:	FLUORESCE	EIN CONJUGA	TION KIT				
Format:	Kit						
Product Type:	Conjugation k	Kit					
Quantity:	1 CONJUGAT	FION for 2mg a	antibody				
roduct Details							
Applications	This product has be	en reported to	work in th	e following applicatio	ons. This information is		
	-			peer-reviewed public	•		
		-		e refer to references i			
	rad-antibodies.com/	-	ecommen	dations, please visit <u>v</u>	<u>www.dio-</u>		
		Yes	No	Not Determined	Suggested Dilution		
	Conjugation	-					
	<b>LYNX Rapid Fluorescein Antibody Conjugation Kit</b> ® enables the rapid conjugation of a pre-prepared lyophilized mixture containing Fluorescein label to an antibody or protein. Activation of proprietary reagents within the antibody-label solution results in directional covalent bonding of Fluorescein to the antibody.						
	The LYNX Rapid Conjugation kit® can be used to label small quantities of antibody/protein at near neutral pH, allowing a high conjugation efficiency with 100% antibody recovery.						
Reagents In The Kit	1 Vial LYNX lyophili: 1 Vial LYNX Modifie		n mix				
	1 Vial LYNX Quench						
Preparing The Antibody		ner reagent					
	The following buffer	-	recommer	nded for preparing the	e antibody:		
	10-50mM amine-free	solutions are e buffer (e.g H	EPES, MI		ohate) pH range 6.5-8.5,		

	<ul> <li>chemicals. Azide (0.02-0.1%), EDTA and common non-buffering salts and sugars have little or no effect on conjugation efficiency.</li> <li>It is recommended that 1-2mg antibody be used in each labelling reaction. For optimal results the antibody should be at a concentration of 1mg/ml, with a maximum volume of 1ml and a recommended antibody amount of 1mg. A maximum of 2mg of antibody can be used to obtain good quality conjugates as long as the maximum conjugation volume of 1ml is not exceeded. Antibody below 1mg/ml can still be used as long as the maximum volume is not exceeded. Using less than the recommended amount of antibody may result in unbound label, but this will be removed during subsequent application wash steps. Antibody below 0.5mg/ml should be concentrated before use with the kit.</li> </ul>
Instructions For Use	1. To the antibody sample add 1ul of the Modifier reagent for every 10ul of antibody and mix gently.
	2. Pipette the mixed antibody-modifier sample directly onto the LYNX lyophilized mix and gently pipette up and down twice to resuspend.
	3. Replace cap onto vial and incubate in the dark at room temperature (20-25 <sup>o</sup> C) for 3 hours or overnight, if preferred.
	4. After incubation, add 1ul of Quencher reagent for every 10ul of antibody used. Leave to stand for 30 minutes before use.
References	1. Walker, E. et al. (2014) Microscopic Detection of Quenched Activity-Based Optical
	Imaging Biol. 16: 608-18.
	<ul> <li>Imaging Probes Using an Antibody Detection System: Localizing Protease Activity. Mol Imaging Biol. 16: 608-18.</li> <li>2. Mahata, B. <i>et al.</i> (2014) Single-cell RNA sequencing reveals T helper cells synthesizing steroids de novo to contribute to immune homeostasis. <u>Cell Rep. 7: 1130-42.</u></li> <li>3. Gawronska-Kozak, B. <i>et al.</i> (2021) Dermal White Adipose Tissue (dWAT) Is Regulated by Foxn1 and Hif-1α during the Early Phase of Skin Wound Healing. <u>Int J Mol Sci. 23 (1):</u></li> </ul>
	<ul> <li>Imaging Probes Using an Antibody Detection System: Localizing Protease Activity. Mol Imaging Biol. 16: 608-18.</li> <li>2. Mahata, B. <i>et al.</i> (2014) Single-cell RNA sequencing reveals T helper cells synthesizing steroids de novo to contribute to immune homeostasis. <u>Cell Rep. 7: 1130-42.</u></li> <li>3. Gawronska-Kozak, B. <i>et al.</i> (2021) Dermal White Adipose Tissue (dWAT) Is Regulated</li> </ul>
Storage	<ul> <li>Imaging Probes Using an Antibody Detection System: Localizing Protease Activity. Mol Imaging Biol. 16: 608-18.</li> <li>2. Mahata, B. <i>et al.</i> (2014) Single-cell RNA sequencing reveals T helper cells synthesizing steroids de novo to contribute to immune homeostasis. <u>Cell Rep. 7: 1130-42.</u></li> <li>3. Gawronska-Kozak, B. <i>et al.</i> (2021) Dermal White Adipose Tissue (dWAT) Is Regulated by Foxn1 and Hif-1α during the Early Phase of Skin Wound Healing. Int J Mol Sci. 23 (1): <u>257.</u></li> <li>4. Jax, E. <i>et al.</i> (2023) Evaluating Effects of AIV Infection Status on Ducks Using a Flow Cytometry-Based Differential Blood Count. <u>Microbiol Spectr. 11 (4): e0435122.</u></li> <li>5. Gawronska-Kozak, B. <i>et al.</i> (2024) Hypoxia and Foxn1 alter the proteomic signature of dermal fibroblasts to redirect scarless wound healing to scar-forming skin wound healing</li> </ul>

Health A Informati	nd Safety ion	https://www.	bio-rad-an Fluorescei agent (105	,		#10549 available at:
Licensec	l Use	Kingdom par purchase of right (withou use the prod	tent numb this produ It the right duct to mal	per 2446088 and assoc uct conveys to the buye to resell repackage or ke conjugates for resea	iated internat er the limited, further sublic arch and deve	patent protected under United ional patent applications. The non exclusive non-transferable cense) under these patents to elopment purposes only. The
		data made u purchasing l at. Endeavo	using this p icenses fo ur House,	product, or its compone or diagnostic and other Langford Business Pa	ents to a third uses may be rk, Langford	r its components, or materials or I party. Further information on e obtained by contacting Bio-Rad Lane, Kidlington, Oxon. OX5 <u>antibodies@bio-rad.com</u>
Regulato	ıry	data made u purchasing l at. Endeavo	using this p icenses fo ur House, D KINGDC	product, or its compone or diagnostic and other Langford Business Pa OM. Tel: +44 1865 852	ents to a third uses may be rk, Langford	l party. Further information on obtained by contacting Bio-Rad Lane, Kidlington, Oxon. OX5
	Tel: +1 800 265 73 Fax: +1 919 878 3 Email: antibody_s	, data made u purchasing I at. Endeavo 1GE UNITEI For research	using this p icenses fo ur House, D KINGDC n purposes Worldwide	product, or its compone or diagnostic and other Langford Business Pa OM. Tel: +44 1865 852	ents to a third uses may be rk, Langford 700. E-mail: Europe	l party. Further information on obtained by contacting Bio-Rad Lane, Kidlington, Oxon. OX5
North & South America	Tel: +1 800 265 73 Fax: +1 919 878 3 Email: antibody_s	, data made u purchasing l at. Endeavou 1GE UNITEI For research For research 76 N 751 ales_us@bio-rad.c	using this p icenses fo ur House, D KINGDC n purposes Worldwide	product, or its compone or diagnostic and other Langford Business Pa OM. Tel: +44 1865 852 s only Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@b	ents to a third uses may be rk, Langford 700. E-mail: Europe	I party. Further information on obtained by contacting Bio-Rad Lane, Kidlington, Oxon. OX5 <u>antibodies@bio-rad.com</u> Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint