

Datasheet: LNK063F BATCH NUMBER 150408

LYNX RAPID	FLUORESCEIN AI	ITIBODY	CONJUGATION	KIT
FLUORESCEI	N CONJUGATION	KIT		
Kit				
Conjugation K	it			
3 CONJUGAT	IONS for 20µg ant	body		
derived from testing communications from information. For gene	within our laboraton the originators. P eral protocol recom protocols.	ies, peer lease refe mendatio	-reviewed publica er to references ir ons, please visit <u>w</u>	ations or personal ndicated for further /ww.bio-
Conjugation		, N	lot Determined	Suggested Dilution
Activation of propriet	ary reagents withir	the antib		• •
•			•	
1 Vial LYNX Modifier		x		
1 Vial LYNX Quench	-			
The following buffer s	er reagent	ımended	for preparing the	antibody:
The following buffer s	er reagent solutions are recor buffer (e.g HEPE	6, MES, N	IOPS and phosp	hate) pH range 6.5-8.5,
	FLUORESCEI Kit Conjugation K 3 CONJUGAT This product has been derived from testing to communications from information. For gene rad-antibodies.com/p Conjugation We recommend that ratio. LYNX Rapid Fluores pre-prepared lyophilit Activation of propriet covalent bonding of the LYNX Rapid Correst antibody/protein at mantibody recovery. 3 Vials LYNX lyophilit	FLUORESCEIN CONJUGATION Kit Conjugation Kit 3 CONJUGATIONS for 20µg antil This product has been reported to work derived from testing within our laborator communications from the originators. Plinformation. For general protocol recommendations rad-antibodies.com/protocols. Yes No Conjugation Ve recommend that for each conjugation We recommend that for eagents within covalent bonding of Fluorescein to the antibody compre-prepared lyophilized mixture contain Activation of proprietary reagents within covalent bonding of Fluorescein to the antibody/protein at near neutral pH, alloantibody recovery.	FLUORESCEIN CONJUGATION KIT Kit Conjugation Kit 3 CONJUGATIONS for 20µg antibody This product has been reported to work in the fold derived from testing within our laboratories, peer communications from the originators. Please reference information. For general protocol recommendation rad-antibodies.com/protocols. Yes No Conjugation • We recommend that for each conjugation the user pre-prepared lyophilized mixture containing Fluo Activation of proprietary reagents within the antilic covalent bonding of Fluorescein to the antibody. The LYNX Rapid Conjugation kit® can be used to antibody/protein at near neutral pH, allowing a hid antibody recovery. 3 Vials LYNX lyophilized Fluorescein mix	Kit Conjugation Kit 3 CONJUGATIONS for 20µg antibody This product has been reported to work in the following application derived from testing within our laboratories, peer-reviewed publicat communications from the originators. Please refer to references in information. For general protocol recommendations, please visit with rad-antibodies.com/protocols. Yes No Not Determined Conjugation • We recommend that for each conjugation the user determines the ratio. LYNX Rapid Fluorescein Antibody Conjugation Kit® enables the pre-prepared lyophilized mixture containing Fluorescein label to a Activation of proprietary reagents within the antibody-label solution covalent bonding of Fluorescein to the antibody. The LYNX Rapid Conjugation kit® can be used to label small qua antibody/protein at near neutral pH, allowing a high conjugation effantibody recovery. 3 Vials LYNX lyophilized Fluorescein mix

	chemicals . Azide (0.02-0.1%), EDTA and common non-buffering salts and sugars have little or no effect on conjugation efficiency.
	It is recommended that 10-20ug antibody be used in each labelling reaction. For optimal results the antibody should be at a concentration of 1mg/ml, with a maximum volume of 10ul and a recommended antibody amount of 10ug. A maximum of 20ug of antibody can be used to obtain good quality conjugates as long as the maximum conjugation volume of 10ul is not exceeded. Antibody below 1mg/ml can still be used as long as the maximum volume is not exceeded. Using less than the recommended amount of antibody may result in unbound label, but this will be removed during subsequent application wash steps. Antibody below 0.5mg/ml should be concentrated before use with the kit.
Instructions For Use	1. To the antibody sample add 1ul of the Modifier reagent for every 10ul of antibody and mix gently.
	2. Pipette the mixed antibody-modifier sample directly onto the LYNX lyophilized mix and gently pipette up and down twice to resuspend.
	3. Replace cap onto vial and incubate in the dark at room temperature (20-25 ^o C) for 3 hours or overnight, if preferred.
	4. After incubation, add 1ul of Quencher reagent for every 10ul of antibody used. Leave to stand for 30 minutes before use.
References	 Walker, E. <i>et al.</i> (2014) Microscopic Detection of Quenched Activity-Based Optical Imaging Probes Using an Antibody Detection System: Localizing Protease Activity. <u>Mol</u> <u>Imaging Biol. 16: 608-18.</u> Mahata, B. <i>et al.</i> (2014) Single-cell RNA sequencing reveals T helper cells synthesizing steroids de novo to contribute to immune homeostasis. <u>Cell Rep. 7: 1130-42.</u>
References Storage	 Imaging Probes Using an Antibody Detection System: Localizing Protease Activity. <u>Mol</u> <u>Imaging Biol. 16: 608-18.</u> 2. Mahata, B. <i>et al.</i> (2014) Single-cell RNA sequencing reveals T helper cells synthesizing
	Imaging Probes Using an Antibody Detection System: Localizing Protease Activity. <u>Mol</u> <u>Imaging Biol. 16: 608-18.</u> 2. Mahata, B. <i>et al.</i> (2014) Single-cell RNA sequencing reveals T helper cells synthesizing steroids de novo to contribute to immune homeostasis. <u>Cell Rep. 7: 1130-42.</u> Store kit at -20°C only. Newly-conjugated antibody can be stored at 4°C. For long term storage however, the addition of a preservative is recommended. Storage in frost-free freezers is not recommended. This product should be stored undiluted.
Storage	Imaging Probes Using an Antibody Detection System: Localizing Protease Activity. <u>Mol</u> <u>Imaging Biol. 16: 608-18.</u> 2. Mahata, B. <i>et al.</i> (2014) Single-cell RNA sequencing reveals T helper cells synthesizing steroids de novo to contribute to immune homeostasis. <u>Cell Rep. 7: 1130-42.</u> Store kit at -20°C only. Newly-conjugated antibody can be stored at 4°C. For long term storage however, the addition of a preservative is recommended. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing.

purchase of this product conveys to the buyer the limited, non exclusive non-transferable right (without the right to resell repackage or further sublicense) under these patents to use the product to make conjugates for research and development purposes only. The purchaser cannot sell or otherwise transfer this product, or its components, or materials or data made using this product, or its components to a third party. Further information on purchasing licenses for diagnostic and other uses may be obtained by contacting Bio-Rad, at. Endeavour House, Langford Business Park, Langford Lane, Kidlington, Oxon. OX5 1GE UNITED KINGDOM. Tel: +44 1865 852 700. E-mail: antibodies@bio-rad.com

Regulatory For research purposes only

North & South Tel: +1 800 265 7376 America Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700 Europe Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M354408:190717'

Printed on 17 Oct 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint