

Datasheet: LNK092PECY5.5

Description:	LYNX RAPID RPE-Cy5.5 ANTIBODY CONJUGATION KIT		
Name:	RPE-Cy5.5 CONJUGATION KIT		
Format:	Kit		
Product Type:	Conjugation Kit		
Quantity:	3 CONJUGATIONS for 60µg antibody		

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol					
	recommendations, please visit www.bio-rad-antibodies.com/protocols.					
	Conjugation	Yes	No	Not Determined	Suggested Dilution	
	Conjugation	_				
	We recommend that for each conjugation the user determines the best antibody:conjugate ratio.					
Product Information	LYNX Rapid RPE-Cy5.5 Antibody Conjugation Kit® enables the rapid conjugation of a pre-prepared lyophilized mixture containing R-Phycoerythrin (RPE)-Cy5.5 label to an antibody o protein. Activation of proprietary reagents within the antibody-label solution results in directional covalent bonding of RPE-Cy5.5 to the antibody.					
	The LYNX Rapid Conjugation kit® can be used to label small quantities of antibody/protein at near neutral pH, allowing a high conjugation efficiency with 100% antibody recovery.					
Reagents In The Kit	3 Vials of 100μg LYNX lyophilized RPE-Cy5.5 mix					
	1 Vial LYNX Modifier reagent					
	1 Vial LYNX Quencher rea	gent				
Preparing The Antibody	The following buffer solutions are recommended for preparing the antibody:					
	10-50mM amine-free buffer (e.g HEPES, MES, MOPS and phosphate) pH range 6.5-8.5, although					
	moderate concentrations of Tris buffer (<20mM) may be tolerated.					
	If possible, avoid buffers containing nucleophilic components such as primary amines and thiols (e.g. thiomersal/thimerosal) since they may react with LYNX chemicals. Azide (0.02-0.1%), EDTA, up to 50% Glycerol and common non-buffering salts and sugars have little or no effect on conjugation efficiency.					
	Due to the large size of the RPE-Cy5.5 label, it is recommended that 50-60ug of antibody be used for every 100ug RPE-Cy5.5, to ensure a slight RPE-Cy5.5 molar excess (50ug antibody gives a 1:1					

for every 100ug RPE-Cy5.5, to ensure a slight RPE-Cy5.5 molar excess (50ug antibody be used hor every 100ug RPE-Cy5.5, to ensure a slight RPE-Cy5.5 molar excess (50ug antibody gives a 1:1 Ab:RPE-Cy5.5 molar ratio). For optimal results the antibody should be at a concentration of 1mg/ml, with a maximum volume of 60ul and a maximum antibody amount of 60ug. Antibody at a concentration of greater than 1mg/ml requires dilution. Antibody below 1mg/ml can still be used as long as the maximum volume is not exceeded. Using less than the recommended amount of antibody may result in unbound label, but this will be removed during subsequent application wash

		steps. Antibody below 0.	5mg/ml should be concentrated before	use with the kit.				
Instructio	ons For Use	1. To the antibody sample add 1μ I of the Modifier reagent for every 10μ I of antibody and mix gently						
		2. Pipette the mixed anti pipette up and down twice	body-modifier sample directly onto the l ce to resuspend.	LYNX lyophilized mix and gently				
		3. Replace cap onto vial overnight if preferred.	and incubate in the dark at room temp	erature (20-25ºC) for 3 hours, or				
		4. After incubation, add for 30 minutes before us	1μl of Quencher reagent for every 10μl e.	of antibody used. Leave to stand				
Storage		preservative is recomme	zers is not recommended. stored undiluted.	storage however, the addition of a				
Guarante	9	12 months from date of	despatch					
Acknowle	edgements		thereof is manufactured under license f per 5,268,486 and related patents. Cy a					
Health An Informatio		Material Safety Datasheet documentation #10540 #10546 #10549 available at: Lyophilized RPE-Cy5.5 Mix (10540): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10540.pdf</u> Modifier Reagent (10546): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10546.pdf</u> Quencher Reagent (10549): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10549.pdf</u>						
Licensed	Use	patent number 2446088 product conveys to the k resell repackage or furth for research and develop product, or its componen third party. Further inform obtained by contacting E	methodology of conjugation are patent and associated international patent app ouyer the limited, non exclusive non-trar er sublicense) under these patents to u oment purposes only. The purchaser can hts, or materials or data made using this nation on purchasing licenses for diagn Bio-Rad, at. Endeavour House, Langford GE UNITED KINGDOM. Tel: +44 1865	blications. The purchase of this nsferable right (without the right to use the product to make conjugates nnot sell or otherwise transfer this s product, or its components to a ostic and other uses may be d Business Park, Langford Lane,				
Regulatory		For research purposes c	nly					
rth & South nerica	Tel: +1 800 265 Fax: +1 919 878 Email: antibody_		Tel: +44 (0)1865 852 700 Europe Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com				
			'M349225:190228'					
			Printed on 20 May 2019					

© 2019 Bio-Rad Laboratories Inc | Legal | Imprint